

FORUM REVIEW ARTICLE

Mitochondria-Associated Membranes: Composition, Molecular Mechanisms, and Physiopathological Implications

Carlotta Giorgi,¹ Sonia Missiroli,¹ Simone Patergnani,¹ Jerzy Duszyński,²
Mariusz R. Wieckowski,^{2,*} and Paolo Pinton^{1,*}

Abstract

Significance: In all cells, the endoplasmic reticulum (ER) and mitochondria are physically connected to form junctions termed mitochondria-associated membranes (MAMs). This subcellular compartment is under intense investigation because it represents a “hot spot” for the intracellular signaling of important pathways, including the synthesis of cholesterol and phospholipids, calcium homeostasis, and reactive oxygen species (ROS) generation and activity. **Recent Advances:** The advanced methods currently used to study this fascinating intracellular microdomain in detail have enabled the identification of the molecular composition of MAMs and their involvement within different physiopathological contexts. **Critical Issues:** Here, we review the knowledge regarding (i) MAMs composition in terms of protein composition, (ii) the relationship between MAMs and ROS, (iii) the involvement of MAMs in cell death programs with particular emphasis within the tumor context, (iv) the emerging role of MAMs during inflammation, and (v) the key role of MAMs alterations in selected neurological disorders. **Future Directions:** Whether alterations in MAMs represent a response to the disease pathogenesis or directly contribute to the disease has not yet been unequivocally established. In any case, the signaling at the MAMs represents a promising pharmacological target for several important human diseases. *Antioxid. Redox Signal.* 22, 995–1019.

Introduction and Mitochondria-Associated Membranes Composition

THE INTRACELLULAR MOBILITY of biological macromolecules and organelles is highly restricted by the high viscosity of the cytosol (87) due to the extreme crowding of intracellular components (46). *In vivo*, nearly all biological macromolecules and other intracellular structures exist, at least transiently, as components of structural and functional complexes (1). This restricted mobility also influences the strength of interactions between various intracellular components. The attractive interactions between components, such as the high affinity of interacting protein structures, represent an additional factor influencing this strength. The direct interactions between mitochondria and the endoplasmic reticulum (ER) within the cell have been extensively

studied both functionally and structurally. The first approach involved studies on the direct exchange of ions and metabolites between the organelles, particularly that of Ca^{2+} ions (142, 154). The close apposition of mitochondria to the ER was found to account for the selective transmission of physiological and pathological Ca^{2+} signals directly from the ER to the mitochondria (60, 145). The second approach involved microscopic visualization of the interactions or isolation of the interacting regions of the organelles and identification of the proteins or lipids present at the sites of interaction. The mutual affinities between these molecules strengthen the interaction between intracellular components, enhancing the half-life of their interaction. When this strength exceeds a certain level and escapes the “kiss and run” formula, it is possible to isolate the interacting fragments of the interacting organelles. Mitochondria-associated membranes (MAMs) (61, 104, 179),

¹Section of Pathology, Oncology and Experimental Biology, Laboratory for Technologies of Advanced Therapies (LTTA), Department of Morphology, Surgery and Experimental Medicine, University of Ferrara, Ferrara, Italy.

²Department of Biochemistry, Nencki Institute of Experimental Biology, Warsaw, Poland.

*These authors share senior co-authorship.

ER-mitochondria encounter structures (96, 97), and plasma membrane-associated membranes (PAMs) (168) represent the best known examples of these preparations.

Among these preparations, MAMs (Figs. 1, 2) are the best characterized. The first reports of the direct association of mitochondria and the ER date from the late 1950s (31), and this type of preparation has been subsequently characterized by numerous groups [for recent reviews, see Refs. (94, 114, 125)]. An isolated MAMs fraction is composed of membrane fragments from both the ER and the outer mitochondrial membrane (OMM) that had been in close contact at the time of cellular subfractionation. More recently, the ER portion of the MAMs fraction has been regarded as a detergent-resistant lipid raft (5, 52).

The contact sites between mitochondria and the ER are dynamic structures that are sensitive to the physiological conditions of the cell. This dynamic nature results in a transient and highly variable MAMs composition. The variety of roles played by the MAMs fraction that have been described is related to their unique lipid and protein composition. The studies performed in the past decade that have identified the molecular components of the MAMs fraction have demonstrated that this fraction may contain numerous proteins [more than 75 according to Raturi and Simmen (149), more than 1000 according to Poston *et al.* (147)]. From this protein composition, it was deduced that MAMs are crucial for numerous cellular processes, including protein sorting [as indicated by the presence of phosphofurin acidic cluster sorting protein 2 (PACS-2)], inflammation [as indicated by the presence of the following inflammasome components: NACHT, LRR and PYD domains-containing protein 3 (NALP3), adaptor ASC and thioredoxin-interacting protein (TXNIP)], ER stress [*e.g.*, 75-kDa glucose-regulated protein (GRP75) and ER resident protein 44 (ERp44)], Ca^{2+} handling [*e.g.*, the inositol 1,4,5-trisphosphate receptor (IP3R), ryanodine receptor, sigma-1 receptor (Sig1R), p53, and promyelocytic leukemia protein (PML)], mitochondrial contact sites [*e.g.*, the voltage-dependent anion channel (VDAC) and adenine nucleotide translocase (ANT)], lipid synthesis and trafficking [*e.g.*, PSS-1 and PSS-2, serine active site containing 1 (SERAC1), FACL1 and FACL4, acyl-CoA desaturase, fatty acid transport protein 4 (FATP4), phosphatidylethanolamine *N*-methyltransferase 2 (PEMT2), and several other proteins; for additional details, (158)], protein folding [*e.g.*, calnexin (CNX)], apoptosis [*e.g.*, Bcl-2 and HCLS1-binding protein 3], and modulation of mitochondrial morphology [mitochondria-shaping pro-

teins and chaperone proteins (mitofusins 1 and 2; MFN1 and MFN2, respectively)]. Moreover, MAMs are the predominant subcellular location for γ -secretase activity and for presenilin-1 (PS1) and presenilin-2 (PS2), two proteins that, when mutated, cause familial Alzheimer's disease (FAD) (4, 5) as discussed later in detail. The ER-mitochondrial cross-talk can also be affected by several viral proteins such as human cytomegalovirus vMIA and the p7 and NS5B proteins of hepatitis C virus, which are also targeted to MAMs. Moreover, it has been found that the tumor suppressors PML and p53 can modulate the ER-mitochondria Ca^{2+} cross-talk by its presence in MAMs (63, 144). In addition, p66Shc, which is a cytosolic adaptor protein that is involved in the cellular response to oxidative stress when phosphorylated at Ser36, has also been identified at the sites of mitochondria-ER association (61, 101). Taking into account that the MAMs fraction contains numerous crucial proteins, as previously mentioned, the involvement of MAMs-mediated disturbances in the pathogenesis of a variety of diseases is not surprising (149). However, the localization of some proteins in the MAMs fraction and the extent of their enrichment remain under debate, and at times, the relationship of these proteins to the MAMs fraction is unclear; see the review by Raturi and Simmen (149).

Relationship Between MAMs and Reactive Oxygen Species

An interesting example of a MAMs-resident reactive oxygen species (ROS)-generating protein is the p66Shc protein. Under physiological conditions, this growth factor adaptor protein (which, in addition to p52Shc and p46Shc, belongs to the ShcA family) is involved in signal transduction *via* the RAS protein. However, under oxidative stress (exogenous and intracellular), p66Shc can participate in the signaling pathway leading to apoptosis. This dual nature of p66Shc arises from the different (tyrosine and serine) phosphorylation sites within this protein. All ShcA proteins possess a similar domain structure; however, p66Shc (which is a product of an alternatively spliced transcript from the SHC1 gene) contains an additional N-terminal proline-rich collagen-homology domain (CH2) containing a serine phosphorylation site (Ser36) that is important for the aforementioned "proapoptotic" properties (40, 170) as well as a functional region (CCB) that is responsible for its interaction with cytochrome *c* (64). As observed for p52Shc and p46Shc, when phosphorylated, p66Shc has been found to bind the





FIG. 1. TEM imaging of MAMs. Representative transmission electron micrographs of mitochondria, ER, and MAMs in mouse embryonic fibroblasts (original magnification, 70,000 \times). The *middle* and *right* panels show a magnification of the selected area. ER, endoplasmic reticulum; MAMs, mitochondria associated membranes. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars

FIG. 2. Determination of MAMs via fluorescence microscopy. HeLa cells were transfected with the SEC61-GFP plasmid to label the ER (green) and loaded with MitoTracker Deep Red to visualize the mitochondrial compartment (red). In the merged and zoomed images, the white areas correspond to the MAMs. Images were obtained using a confocal laser scanning microscope at $63\times$ magnification. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars



Grb2/SOS complex and, in this way, competes with p52Shc for binding to Grb2. This finding suggests that p66Shc can function as a dominant negative regulator of the RAS-mediated signaling pathway (122, 131). Under oxidative stress caused by UV radiation or H_2O_2 treatment, p66Shc can be phosphorylated at Ser36 by one of the serine-threonine kinases [protein kinase $C\beta$ (PKC β) (59), c-Jun N-terminal kinase (JNK), or extracellular signal-regulated kinase (ERK)] (79, 100, 146). Moreover, phosphorylation of p66Shc at Ser36 can be mediated by apoptosis signal-regulating kinase 1 (ASK1)-JNK phosphorylation (110). This phosphorylation initiates the following cascade of events: (i) isomerization of Ser36-phosphorylated p66Shc by the prolyl isomerase Pin1, (ii) dephosphorylation of Ser36-P-p66Shc by phosphatase A2 (PP2A), (10) and (iii) translocation of p66Shc to the mitochondria and/or MAMs fraction, where it participates in ROS production (64, 146) (Fig. 3). Additional details on the “nature” of p66Shc can be found in the comprehensive review by Migliaccio *et al.* (121). It should also be noted that the ROS-sensitive fluorescent probes used in studies describing the ability of MAMs-resident proteins to produce different ROS unfortunately cannot precisely detect ROS. For example, the *CM-H₂DCF* probe used for the measurement of hydrogen peroxide can also detect other ROS, due to

the fact that the oxidation of H_2DCF to DCF is a two-step process: First, the DCF radical is formed, and it is then (second step) oxidized to DCF in a reaction with molecular oxygen (163, 176). Unfortunately, the first step of H_2DCF oxidation can be mediated by various radical species, such as hydroxyl radicals, carbonate radicals, and nitrogen dioxide, as well as by thiyl radicals resulting from thiol oxidation (42, 71, 88, 181, 182). Moreover, alteration of the fluorescent probe signal can be caused by superoxide radicals (formed in the second step of H_2DCF oxidation), which can be dismutated to hydrogen peroxide and cause self-amplification of the signal (67, 128). Similarly, *dihydroethidium* (DHE), and MitoSOX, which are used for the measurement of superoxide, are also not completely specific. For example, DHE and MitoSOX can undergo unspecific oxidation by $ONOO^-$ or $\bullet OH$ into ethidium or mito-ethidium, respectively (88). For this reason, even if the exact form of ROS (*e.g.*, superoxide or H_2O_2) is stated in the text of the original paper commented on in our review, other ROS should also be taken into consideration due to the constraints imposed by probe chemistry.

The first reports addressing the cytosolic localization of p66Shc indicated that it could also be localized in the mitochondrial matrix. Moreover, Orsini *et al.* found that p66Shc can interact with mitochondrial Hsp70 and regulate Ψ_m




FIG. 3. A summary of the MAM-enriched proteins involved in ROS production. During hypoxic condition, the ER-MAM-resident proteins Ero1- $L\alpha$ and ERp44 interact and modulate ER-mitochondria Ca^{2+} communication. In addition, elevated levels of Ero1- $L\alpha$ may promote activation of the UPR-system with a consequent increase in ROS production and Ca^{2+} release from ER. Another MAM protein, p66Shc, is activated under oxidative stress. In parallel, p66Shc can translocate to the mitochondria, where it participates in ROS formation. ERK, extracellular-signal-regulated kinase; JNK, Jun amino-terminal kinases; PP2A, protein phosphatase 2A; Pin1, peptidyl-prolyl cis/trans isomerase; ROS, reactive oxygen species; cyt.c, cytochrome c; Ero1 α , oxidoreductase endoplasmic reticulum oxidoreductin-1 alpha; ERp44, endoplasmic reticulum resident protein 44; UPR, unfolded protein response. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars

(mitochondrial membrane potential) (133). Subsequent studies have revealed that mitochondrial p66Shc is present in the mitochondrial intermembrane space (IMS), where it interacts with cytochrome *c* and, as a redox enzyme, produces H_2O_2 (64). However, the determination that nearly 35% of the mitochondrial p66Shc localizes within the IMS (64) appears to be an overestimation (101, 179). Regarding this model, Gertz *et al.* have proposed that the N-terminus of p66Shc forms a redox module and serves as a thiol-based redox sensor. Under oxidizing conditions, p66Shc can be “activated” due to the formation of two disulfide bonds, resulting in the reversible tetramerization of p66Shc. In contrast, under normal conditions, glutathione and thioredoxins protect against p66Shc tetramerization by reducing potentially formed disulfide bonds and, in this way, inactivating the proapoptotic properties of p66Shc (56).

Direct involvement of p66Shc in mitochondrial ROS production has been repeatedly described by several groups (37, 64, 102, 103, 146). However, it remains unclear whether p66Shc can translocate across the OMM and resides in the IMS, where it can interact with cytochrome *c* (64), or whether it binds to the OMM from the cytosolic side (101). In the model in which p66Shc is localized in the IMS, p66Shc has been proposed to interact with cytochrome *c* (due to the CCB domain present in p66Shc) and transfers electrons from cytochrome *c* to molecular oxygen, which results in H_2O_2 formation. p66Shc present in the MAMs fraction can interact with an as yet unidentified OMM protein and, thus, can participate in ROS production (Fig. 3). Our studies have shown that p66Shc can be found not only in the cytosol and MAMs (179) but also in the PAM fraction (101). Interestingly, the level of p66Shc in the MAMs fraction increases in an age-dependent manner, corresponding well to mitochondrial H_2O_2 production, which

has been found to increase with age (101). Lebedzinska *et al.* performed studies on fibroblasts from patients with diagnosed mitochondrial disorders and demonstrated that p66Shc can also be phosphorylated at Ser36 in the event of intracellular oxidative stress caused by mitochondrial dysfunction (102, 103). In the case of these studies, it is very difficult to distinguish whether the “activated” p66Shc pathway in fibroblasts from patients participates in superoxide ($O_2^{\cdot-}$) or H_2O_2 production. Admittedly, the administration of hispidin (an inhibitor of PKC β phosphorylation of p66Shc) decreases cytosolic and mitochondrial $O_2^{\cdot-}$ levels in these cells, indicating that p66Shc may be involved in the production of this type of ROS. However, hispidin also causes an increase in the levels of cytosolic and mitochondrial dismutases (SOD1 and SOD2), which are enzymes catalyzing the reduction of superoxide anions to H_2O_2 . Therefore, a decrease in the $O_2^{\cdot-}$ level after hispidin treatment not only can be caused by the inhibition of p66Shc Ser36 phosphorylation but also may be SOD1 and SOD2 specific. ROS production by p66Shc appears to be a specialized function whereby electrons are removed from the ETC to catalyze the partial reduction of molecular oxygen (64). The redox activity of p66Shc accounts for the decrease in ROS levels observed in p66Shc knockout cells (121) and is also responsible for an altered mitochondrial metabolism under basal conditions that is characterized by lower oxygen consumption (129).

Interestingly, the connection between p66Shc and ROS seems to have significant physiological relevance in the case of hypertension. It appears that p66shc mediates hypertension-associated, cyclic stretch-dependent, endothelial damage and that in stressed cells, activation of integrin $\alpha 5\beta 1$ and c-Jun N-terminal kinase enhances the phosphorylation p66shc at Ser36 and, thus, ROS production (166). The results of these

experiments can be extrapolated to an organismal level as well. It has been demonstrated that in mice lacking p66shc, age-related and hyperglycemia-induced endothelial dysfunctions (24, 32, 65, 93) as well as the extent of atherosclerosis (126) are diminished.

Additional examples of proteins that can produce ROS and that are localized in the MAMs fraction include Ero1- $L\alpha$ (which is present in several tissues and most cell types) and Ero1- $L\beta$ (which is abundant in cells with a high secretory capacity) (2, 47, 149). Although Ero1- $L\alpha$ is a luminal ER oxidoreductase, it presumably binds to the ER membranes in regions involved in MAMs formation, resulting in greater than 75% of Ero1- $L\alpha$ localization in the MAMs fraction (58). Ero1- $L\alpha$, in addition to protein disulfide-isomerase (PDI), is responsible for the formation of disulfide bonds and, hence, plays an essential role in protein folding (21, 45). Enyedi *et al.* have shown that the activity of Ero1- $L\alpha$ results in significant H₂O₂ formation in the ER. Briefly, oxidative protein folding consists of two major steps: (i) The FAD-bound Ero1 protein oxidizes the PDI (*e.g.*, ERp44 or ERp57) and (ii) PDI subsequently catalyzes the formation of disulfide bonds within newly synthesized/folding proteins (3). Ero1- $L\alpha$ uses oxygen, which is finally reduced to H₂O₂ as the electron acceptor on the oxidation of PDI (74). Studies performed on yeast have indicated that greater than 25% of the ROS produced during protein synthesis/folding is related to yeast homolog Ero1p activity (172). The interplay between Ero1- $L\alpha$ and ERp44, which is an ER luminal chaperone protein (with thioredoxin activity) that is also present in the MAMs fraction, is responsible for the regulation of Ca²⁺ release from the ER *via* IP3R1. Ero1- $L\alpha$ in the ER-mitochondrial hot spots interacts with IP3R1, oxidizing it to potentiate the release of Ca²⁺ during ER stress (2). Interestingly, hypoxic conditions result in the complete relocation of Ero1- $L\alpha$ from MAMs, indicating that its intracellular localization depends on oxidizing conditions (58). Moreover, the ability of Ero1- $L\alpha$ to modulate ER-mitochondria Ca²⁺ communication may substantially affect the induction of apoptosis (40, 169). In contrast, ERp44 interactions with IP3R1 under reducing ER conditions inhibit Ca²⁺ transfer to the mitochondria (76); however, the oxidation of IP3R1 by Ero1- $L\alpha$ is accompanied by ERp44 dissociation from the complex with IP3R1, which enables its full activation (2, 109) (Fig. 3).

Additional details on these topics can be found in the following reviews (149, 165).

MAMs in Cell Death Pathways

Several studies have highlighted and broadened the functional roles of MAMs in a variety of cellular processes, from lipid synthesis/transport, Ca²⁺-signaling, and ER stress to mitochondrial morphology.

In addition to these roles, MAMs play a key role in the initiation and amplification of cell death. In the next sections, we will illustrate how these contact sites serve as signaling platforms that are capable of determining cellular life and death decisions through the regulation of programmed cell death events.

Apoptosis

In the mitochondrial pathway of apoptosis, stress signals induce mitochondrial outer membrane permeabilization

(MMP), which results in permeabilization of the OMM. MMP then facilitates the release of several proteins that usually reside in the mitochondrial intermembrane and intracristal space. These proteins include cytochrome C (cyt-C), apoptosis-inducing factor (AIF), and Smac/DIABLO. Finally, these proteins trigger or facilitate the formation of a caspase-activating complex, the apoptosome, which results in the activation of effector caspases (66). It is therefore clear that one of the key steps of mitochondrial apoptosis is the permeabilization of the OMM. This event is highly regulated by members of the BCL-2 family, particularly through the activity of BAX, BAK, and BID. Under normal conditions, the proapoptotic protein BAK is situated in the cytosolic compartment. On apoptosis induction, BAX inserts into the OMM, where it associates with BAK or BID. Such complexes stimulate MMP and the release of cyt-C by forming pores in the OMM. Nevertheless, how this molecular opening induced by BAX/BAK/BID occurs is still controversial. A first putative mechanism suggested the cooperation of BAX with other proteins to form and open a pore in the inner membrane (the so-called mitochondrial permeability transition pore [mPTP], permeability transition pore complex) allowing water and proteins till ~ 1.5 kDa to pass through (98).

The mPTP is a multiprotein complex that forms at the junctions between the inner and outer mitochondrial membranes. The current mPTP model is built around the F1/Fo ATP-synthase and is composed of several elements, including ANT, VDAC, BAX, and BAK (12, 14, 15).

Opening the pore results in matrix swelling and OMM disruption, which, in turn, promote the release of proteins from the IMS.

An alternative model of MMP was proposed, which did not indicate a key role for the PTP. This mechanism is regulated by BCL-2 proteins, which act directly on the OMM. For example, it was found that anti-apoptotic BCL-2 family members work to block the MMP, while proapoptotic members can act to activate BAK/BAX/BID or interfere with anti-apoptotic BCL-2 family members (48, 99).

Other studies suggest a critical role of VDAC in the MMP and apoptosis (162). Initially, the overexpression of VDAC leads to apoptosis in a variety of cell types (186). In addition, several works address the molecular pathway involving VDAC. In fact, under physiological conditions, antiapoptotic BCL-2 members interact with VDAC and regulate its function, which is to shuttle ATP from the mitochondrial matrix to the cytoplasm (73). However, the scenario is not actually quite so simple: Recent evidence shows that the contribution of VDAC to cell death can be isoform and stimulus dependent. Accordingly, it has been found that VDAC1 silencing promotes apoptosis, whereas silencing of VDAC2 has the opposite effect (38).

MMP is not the only crucial event in apoptosis. There is also a major change in the membrane potential (in the plasma membrane potential [Ψ_{pm}] and the mitochondrial transmembrane potential [Ψ_m]) accompanying MMP. Under normal conditions, the K⁺ concentration is much higher in the cytosol than in the extracellular fluid. A continuous, low K⁺ efflux, *via* K⁺-channels, is essential for the maintenance of Ψ_{pm} , which is vital for ion and volume homeostasis. It has been reported that during mitochondrial swelling and MMP, Ψ_{pm} collapses, resulting in dissipation of intracellular [K⁺], cell shrinkage, DNA fragmentation, and loss of membrane

asymmetry, with aberrant exposure of phosphatidylserine residues on the plasma membrane surface (35). Thus, a gross perturbation of Ψ_{pm} appears to occur in the postmitochondrial stage of apoptosis.

Ψ_m also contributes to mitochondrial apoptosis. A prominent dissipation of Ψ_m takes place in the early stage of the apoptotic process after the opening of the PTP. Nevertheless, studies suggest that loss of Ψ_m could be a consequence of the apoptotic-signaling pathway. For example, during the etoposide-induced apoptosis of L929 fibroblasts, loss of Ψ_m occurs as a late event after nuclear alterations and BAX translocation to the mitochondria (91). Moreover, it has been suggested that loss of Ψ_m is not required for cyt-C release, but only for the release of AIF (112). Considering these remarks, it is clear that dissipation of Ψ_m is a classic feature of apoptosis, but its effective role in this pathway remains to be addressed.

Cardiolipin. To play crucial roles in cellular bioenergetics and cell survival, mitochondria require the correct import of a large number of proteins from the cytosol. Lipids play a key role in this “mitochondrial protein sorting.” Of the mitochondrial lipids, the dimeric phospholipid CL is characteristic of this organelle, and it is responsible for the stability of several IMM protein complexes (33). Reduced levels of CL induce a collapse of Ψ_m with consequent blocking of the Ψ_m -

dependent protein translocation into mitochondria. Moreover, during apoptosis, CL is redistributed between the IMM and the OMM, and its oxidation induces the release of proapoptotic factors (53, 86). Although the molecular mechanism of CL translocation remains elusive, MAMs have been proposed to play a key role in CL movement. In demonstration of this theory, the state and efficiency of MAMs are prime determinants for CL transfer and recruitment to the OMM (78, 152) (Fig. 4).

Mitochondrial network. The shape of the mitochondrial network is also important to the functioning and health of the cell. During apoptosis, the mitochondrial network undergoes dramatic rearrangements, and it is generally accepted that it collapses into small spherical structures in response to apoptotic stimuli.

Furthermore, as discussed earlier, mitochondrial dynamics and morphology are modulated by a correct assemblage of ER-mitochondria contacts. In wild-type cells, during apoptosis, mitochondrial fission facilitates mitochondrial fragmentation, the collapse of Ψ_m , and the release of IMS-stored apoptogenic factors. For example, inhibition of fission through DRP1 RNA interference retards the release of cyt-C from the IMS (107). Consistent with these data, DRP1-KO-derived MEFs display a delay in cyt-C release and retardation of caspase activation (81), and RNA interference targeting




FIG. 4. A schematic overview of the main MAM-associated proteins involved in apoptosis. A growing body of evidence highlights the importance of several MAM-related proteins in the regulation of the apoptotic pathway. Some of these proteins can modulate apoptosis through the generation of ROS, formation of cardiolipin (CL) microdomains, or recruitment and regulation of fission/fusion proteins. Other proteins that link the ER to the mitochondrial surface (particularly PML, AKT, PTEN and RAS) are able to modulate the execution of apoptosis by controlling Ca^{2+} waves in stressed cells. As a result, excess Ca^{2+} flowing out of the ER is taken up by the mitochondria (through the uniporter complex), promoting the execution of the apoptotic program. DRP1, dynamin-related protein 1; MFN2, mitofusin 2; PACS2, phosphofurin acidic cluster sorting protein 2; PERK1, protein kinase RNA-like endoplasmic reticulum kinase; IRE1- α , inositol-requiring kinase 1; ATF6, activating transcription factor 6; FIS1, fission 1 (mitochondrial outer membrane) homolog (*S. cerevisiae*); BAP31, mammalian B-cell receptor-associated protein 31; HK2, hexokinase2; cyPd, cyclophilin-D; PP2A, protein phosphatase 2A; ERO1- α , oxidoreductase endoplasmic reticulum oxidoreductin-1 alpha; ERP44, endoplasmic reticulum resident protein 44. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars

MFF induces mitochondrial elongation, with a consequent delay in cyt-C release and blocking of the apoptotic program.

Taken together, these data suggest an important role of fission-related proteins during the progression of the apoptotic program. After the induction of apoptosis, the fission protein DRP1 translocates to the OMM, with consequent augmentation of mitochondrial fragmentation. Indeed, it has been reported that during apoptosis cytosolic BAX/BAK proteins travel to the OMM and interact with DRP1 and MFN2. As a result, these BCL-2 member proteins promote SUMO modifications of DRP1, which stably associates with the OMM (18, 177). In addition, BAX/BAK induce tBID activation, which is correlated with blocking of mitochondrial fusion, most likely through inhibition of the mitofusin protein MFN-2 (90). The activation of fusion proteins (such as MFN-1/-2 and OPA1) that are indispensable for the maintenance of normal mitochondrial morphology and calcium uptake by mitochondria antagonizes apoptosis progression (51).

Accordingly, inhibition of OPA1-mediated fusion through RNA interference leads to fragmentation of the mitochondrial network, concomitant with the dissipation of the mitochondrial membrane potential. As a consequence of these events, cyt-C is released, and executioner caspases are activated (28, 132).

The apoptotic program is also promoted by depletion of PACS2. Indeed, the absence of PACS-2 induces the caspase-dependent cleavage of BAP31 to yield the pro-apoptotic fragment p20, causing mitochondria to fragment and uncouple from the ER (164).

Overall, these findings indicate that disruption of the apposition of mitochondria with the ER and the consequent increase in mitochondrial fragmentation are functionally linked to apoptosis induction.

Calcium (Ca^{2+}). Ca^{2+} homeostasis is fundamental to numerous cellular mechanisms, including cell death. Elevation of the intracellular Ca^{2+} concentration is dependent on either Ca^{2+} influx from the extracellular space through the plasma membrane or Ca^{2+} release from intracellular Ca^{2+} stores, such as those in the ER. The mitochondria are equally important for Ca^{2+} signaling. Different works have demonstrated that Ca^{2+} release from the ER results in cytosolic Ca^{2+} increases that are paralleled by similar or even increased cycles of mitochondrial Ca^{2+} uptake (11, 116).

This unique characteristic of the mitochondria is principally due to the large electrochemical gradient Ψ_m and due to the existence of MAMs. As describe earlier, the close apposition of the mitochondria and ER creates Ca^{2+} hotspots, which have been found to play pivotal roles in several cellular functions, including the highly efficient transmission of Ca^{2+} from the ER to the adjacent mitochondrial network to stimulate oxidative metabolism. Apoptosis is also intimately connected to the regulation of Ca^{2+} handling promoted by MAMs. An excess of Ca^{2+} flows out of the ER into mitochondria *via* IP₃ receptors (which present at high levels in the MAMs compartment) and promotes the apoptotic program (142). For example, several studies have correlated MAMs with increased Ca^{2+} transfer in sensitization to apoptosis. For example, the Ca^{2+} fluxes into mitochondria induce the oligomerization and activation of BAX, which promotes the permeabilization of the OMM and, ultimately, the release of

pro-apoptotic factors into the cytosol. Other proteins that link the ER to mitochondria have been found to exert control over pro-apoptotic Ca^{2+} fluxes in stressed cells.

A growing body of research highlights the key role of the oncogene H-RAS in the maintenance of tumor survival and proliferation. Although the clear molecular mechanisms underlying these processes are not well established, a recent study illustrated a direct link between Ca^{2+} regulation and HRAS-driven transformation. In this work, Rimessi *et al.* identified HRAS localized to the MAMs compartment, suggesting the possibility that this localization may serve as a strategic point to regulate the transmission of Ca^{2+} from the ER to mitochondria. Indeed, after the induction of oncogenic HRAS, global intracellular Ca^{2+} perturbation accompanied by a dysfunction of mitochondrial physiology has been observed (153).

PTEN (protein phosphatase and tensin homolog deleted on chromosome 10), a tumor suppressor that downregulates AKT and localizes to the MAMs compartment, was recently shown to be able to interact with the AKT/IP3R complex (11), leading to a reduction of its phosphorylation and an increase in Ca^{2+} release. The tumor suppressor PML also modulates the ER-mitochondria Ca^{2+} signaling platform. This protein was found to localize to the ER and MAMs, where it interacts with IP3R3, regulating the ER-mitochondria Ca^{2+} flux and apoptosis. Furthermore, using PML-KO-derived MEFs and a PML chimera that exclusively localizes to the outer surface of the ER (erPML), our laboratory demonstrated that PML also acts as a suppressor of other oncogenic pathways. Indeed, loss of PML promotes a reduction of PP2A activity at the ER and an increase of AKT activity. Consequently, the hyper-phosphorylation of IP3Rs mediated by AKT inhibits ER Ca^{2+} release and causes the cell to be less sensitive to Ca^{2+} -mediated apoptotic stimulation (62).

Other central players in the ER-mitochondria Ca^{2+} flux include a series of chaperones and oxidoreductase, which also localize to the ER/MAMs compartment. ERp44, an ER luminal protein of the thioredoxin family, was found to directly interact with IP3R1, inhibiting its channel activity. Specifically, Higo *et al.* demonstrated that ERp44 directly inhibits the channel activity of IP3R1 in a pH-, redox state-, and $[\text{Ca}^{2+}]_{\text{ER}}$ -dependent manner (76). Furthermore, in this study, it was demonstrated that ERp44 also reinforces the ERO1 α /oxidoreductase system. ERO1- α is an enzyme that modulates the activity of PDI. ERO1- α localizes to the MAMs and ER, where it can interact with and control the activity of HIF1- α , modulating the hypoxic response to disulfide bond formation (120). Based on these observations, the reported intimate relationship between ERp44 and ERO1- α could play a key role in the modulation of IP3R-dependent calcium signaling and, thus, in the regulation of the apoptotic program.

It is clear that the release of Ca^{2+} from ER stores is an essential component of cell survival processes, particularly apoptosis. On release from the ER, Ca^{2+} is taken up by mitochondria through the mitochondrial calcium channel uniporter (MCU), which together with its regulators (MICU1, MICU2, MCUB, EMRE, MCUR1, and miR-25), it constitutes the MCU complex (113, 115). Next, the Ca^{2+} flux promotes apoptosis induction through the activation of the mPTP, which, in turn, promotes cyt-C release and caspase activation. As described earlier, the mPTP is composed of

several components, including proteins localized at MAMs (12). Based on this observation and considering the key role of the mPTP in cell survival, it is possible to predict a direct and novel role of ER-mitochondria apposition in mPTP regulation and, thus, in the onset of apoptosis. Several findings suggest the existence of this intimate relationship. mPTP opening is an important event in cardiomyocyte cell death during ischemia-reperfusion (I/R). In addition, peroxidation of CL by ROS in the presence of Ca^{2+} induces mitochondrial permeability transition and cyt-C release in rat heart mitochondria. Thus, increased levels of peroxidized CL and Ca^{2+} might lead to the opening of the mPTP. Furthermore, these effects were not observed for nonoxidized CL and were inhibited by cyclosporin A and bongkrekic acid (138, 143). Considering that MAMs are prime determinants of CL homeostasis and Ca^{2+} handling, it is possible that these membranes play an active role in the regulation of mPTP opening during I/R.

Other indications of the existence of this relationship might be found by considering p66Shc. As described earlier, p66Shc is a MAMs protein that is strongly regulated by ROS and is involved in Ca^{2+} handling. Once activated, p66Shc oxidizes reduced cyt-C and catalyzes the reduction of O_2 to H_2O_2 using electrons from the respiratory chain. The generated H_2O_2 leads to mPTP opening and consequent caspase activation and apoptosis (64).

The ER not only is one of the main sites of intracellular Ca^{2+} storage in the cell but also is considered the primary site for the synthesis and folding of proteins for the entire cell. When the capacity of the ER machinery to fold proteins becomes inadequate, the cell enters into a dangerous state, known as "ER stress." When ER stress becomes too severe and the dedicated signaling pathways (such as UPR) cannot alleviate this condition, a lethal signal may be turned on, giving rise to cell death, usually in form of apoptosis.

On ER stress three sensor proteins of UPR-system (ATF6, PERK, and IRE1- α) generally stimulate an increase of ER and a gain of function of chaperone activity. Consequently, the ER morphology considerably undergoes changes, and the ER-mitochondria contact sites become stronger and closer. This increased coupling results in major oxygen consumption, a stronger reductive capacity, and augmented Ca^{2+} transfer. This increase in mitochondrial activity is observed only where the reticular and mitochondrial networks are redistributed. Overall, these findings suggest that the increased juxtaposition between the ER and mitochondria observed during ER stress could contribute favorably to cellular adaptation to stressful conditions (17).

Nevertheless, these dependent ER stress sensor proteins can modulate the transmission of signals between the ER and mitochondria in a UPR-independent manner. For example, ablation of the protein PERK, which was recently discovered to be enriched in the MAMs fraction, in addition to the ER compartment, promotes perturbations of ER-mitochondria contact sites and reduces apoptotic activity in response to agents inducing ROS production, Ca^{2+} transfer, and ER stress. Collectively, these data reveal that a conserved MAMs structure is indispensable for transmitting Ca^{2+} as well as ROS-mediated signals to the mitochondria after ROS-based ER stress (174). Accordingly, cells lacking MFN2 exposed to ER stress display a weaker ER-mitochondria interaction and a reduction of apoptosis. Furthermore, other recent studies

have highlighted the key role of the ER-mitochondria juxtaposition in propagating apoptosis in the presence of pro-oxidant inducers of ER stress (25, 62, 137).

Autophagy

MAMs are not only important for the coordination of mitochondrial integrity, Ca^{2+} handling, and apoptotic activity but also a primary element for the initiation and execution of the autophagic machinery. The mitochondria and ER are involved in autophagosome biogenesis. Initially, several ER proteins were found to localize to the autophagosome membrane. Then, electron and fluorescent microscopy analyses demonstrated that the ER and the initial isolation membrane of autophagosomes (the phagophore) are juxtaposed, and on starvation-induced autophagy the cell forms the so-called omegasome, a phosphatidylinositol 3-phosphate compartment connected to the ER that is fundamental for autophagosome formation. Alternatively, the autophagosome might be derived from mitochondria. It has been demonstrated that during starvation, the OMM "gifts" mitochondrial-derived membranes that will be used for autophagosome biogenesis. Furthermore, a crucial role of the ER-mitochondria connection in autophagy induction has been demonstrated. Indeed, MFN2 KO-derived cells are not able to undergo autophagy, supporting the possibility of a contribution of lipids from the ER to mitochondria (70). In addition, these studies provided a further indication of the intimate relationship between MAMs and the autophagosome. However, the importance of ER-mitochondria contact sites for autophagosome formation remained controversial.

This question was answered in 2013 by an elegant study published in *Nature*, where it was demonstrated that autophagosome formation starts at MAMs. Immunofluorescence and electron microscopy, in addition to subcellular fractionation, showed that specific preautophagosome/autophagosome markers (ATG14 and ATG5) localize to MAMs under starved conditions. Consistent with these findings, on starvation, DCFP1 (double FYVE domain-containing protein 1) translocates to the omegasome. In contrast, in PACS2 KO and MFN2 KO cells, both the accumulation of autophagic markers and the translocation of ER-related proteins are significantly reduced, indicating a stable role of MAMs during the completion of autophagosomes (72). In spite of these findings, the role of MAMs in the molecular mechanism of autophagy remains to be elucidated.

It is well known that the main regulator of autophagy is the serine/threonine kinase mTOR (mechanistic target of rapamycin) (95). This kinase exists in two protein complexes: mTOR complex-1 (mTORC1) and -2 (mTORC2). Interestingly, it has been found that mTORC2 localizes to MAMs, where it activates AKT. Once activated, AKT appears to control MAMs integrity and mitochondrial physiology through PACS and hexokinase (HK2) phosphorylation. Furthermore, mTORC2-AKT regulates IP3R3 phosphorylation and Ca^{2+} release at MAMs (8). Despite this, the involvement of mTORC-2 in autophagy remains debatable. Nevertheless, it has been found that Rab32 (described earlier to be an MAMs protein essential for autophagosome formation) is critical for the regulation of mTORC2 activity (29). Overall, these findings may suggest an active role for this

complex in autophagosome formation and the regulation of autophagic activity.

Other MAMs proteins also appear to regulate autophagy. One example is p66Shc, a protein that plays an important role in controlling the mammalian life span (146). It has recently been found that PKC β and p66Shc overexpression leads to a reduction of autophagic activity and their roles in the modulation of autophagy appear to be interrelated. In support of this hypothesis, overexpression of PKC β drives a strong increment of p66Shc phosphorylation and an augmentation of the transfer of p66Shc to the mitochondrial compartment. Consistent with this finding, MEFs derived from PKC β KO mice display a strong and significant reduction of both the phosphorylation and mitochondrial localization of p66Shc (141).

As has been well reviewed, different forms of specialized autophagy have been discovered in recent years (75). One of the most important is mitophagy, which is responsible for the selective removal of damaged and exhausted mitochondria. Several studies have demonstrated that two genes, *PINK1* and *PARK2*, are involved in the maintenance of a healthy population of mitochondria.

Parkin, encoded by the *PARK2* gene is a cytosolic E3 ubiquitin ligase that mediates the ubiquitylation of a number of target proteins. Parkin has been found to be cytoprotective under various conditions, and it has been reported to play a role in mitochondria under stress conditions (127). Parkin ubiquitinates several OMM proteins, including VDAC, MFN, DRP1, BCL2, and BAX (54, 55). When severe damage impacts a population of mitochondria, the PINK1-parkin axis recognizes defective mitochondria, rapidly isolates them from the mitochondrial network, and, finally, degrades them through the ubiquitin-proteasome and autophagic pathways (151). Recently, it has been demonstrated that mitochondria

supply membrane material not only during serum starvation but also during drug-induced autophagy, introducing a novel mechanism of parkin-associated mitophagy.

Cook *et al.* demonstrated that stress and starvation conditions and treatment with drugs that are capable of promoting autophagy increased both PINK1 and parkin localization to the mitochondrial compartment. Furthermore, using confocal and electron microscopy, these authors showed that mitochondria labeled with parkin are not only engulfed by the forming autophagosome but also used to form new a autophagosome. Accordingly, inhibition of the mitophagic process through PINK1 knockdown restores normal functional mitophagy in cells (30). In conclusion, this work could open new avenues for the discovery of novel roles of the ER-mitochondria contact sites in the regulation of the molecular pathways of general and selective autophagy (Fig. 5).

MAMs and Inflammation

In addition to their established role as a signaling hub for Ca²⁺ and lipid transfer between the ER and mitochondria, MAMs have recently been shown to play a central role in the modulation of various key processes, including inflammasome signaling. A link between inflammation and the ER-mitochondria interface was established for the first time in 2011 in a study by Zhou *et al.*, who demonstrated a new role for mitochondria in NLRP3 inflammasome activation (190).

The NOD-like receptors (NLRs) are composed of 22 human genes that are characterized by the presence of a central nucleotide-binding oligomerization (NACHT) domain, C-terminal leucine-rich repeats (LRRs), and an N-terminal effector domain. On activation, select NLR family members form multiprotein complexes (termed inflammasomes) that




FIG. 5. The importance of ER-mitochondria contact sites in autophagy induction and execution. Although the biogenesis of the autophagosome is complex and incompletely understood, it has been demonstrated that MAMs contribute to the initiation of the isolation membrane of the autophagosome. For example, several autophagy-related protein reside in MAM compartments, and MAM proteins are also critical regulators of the execution of the autophagic machinery. For instance, on activation by PKC β (protein kinase C- β), the MAM protein p66Shc promotes autophagy. Furthermore, a recent study highlighting the critical role of mitochondria in supplying membrane material during the mitophagic mechanism could open new avenues for the investigation of novel roles of MAMs in the regulation of the molecular pathway of selective autophagy. ATG, autophagy-related; DFCP1, double FYVE domain-containing protein 1; Rab32, small GTP-binding protein 32 of the RAB family; PINK1, PTEN-induced putative kinase 1. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars

serve as platforms for caspase-1 activation and the subsequent proteolytic maturation of the potent proinflammatory cytokine IL-1 β . (171). Due to its association with numerous inflammatory diseases, the NLRP3 inflammasome is currently the most fully characterized, well-studied inflammasome. The key components of a functional NLRP3 inflammasome include NLRP3, the adaptor protein ASC (apoptosis-associated speck-like protein containing a caspase recruitment domain), and caspase-1 (159). On sensing a wide variety of danger signals, NLRP3 oligomerization and recruitment of ASC and procaspase-1 trigger the autoactivation of caspase-1 and the maturation and secretion of proinflammatory cytokines, such as IL-1 β .

Resting NLRP3 localizes to the cytosol and ER structures, whereas on inflammasome activation using nigericin (an antibiotic) or monosodium urate, both NLRP3 and its adaptor ASC colocalize to the MAMs fraction. Thus, by virtue of its ER-mitochondria localization on activation, the NLRP3 inflammasome is strategically located to receive signals emanating from mitochondria.

Previous studies have indicated that ROS represent a common integrator across several stimuli that activate the NLRP3 inflammasome. In addition to the ER and peroxisomes (19), mitochondria are the primary source of ROS. However, the source of NLRP3-activating ROS and the associated underlying mechanisms remain unclear.

Zhou and colleagues observed a correlation between mitochondrial ROS activity and the presence of IL-1 β in the supernatant of the human THP1 macrophage cell line. To avoid cellular damage, ROS-generating mitochondria

are constantly removed by mitophagy; these authors found that inhibition of the autophagic machinery affects IL-1 β production, resulting in the accumulation of defective and ROS-producing mitochondria, which potentiates NLRP3-dependent inflammasome activation.

Although these results indicated that the prolonged presence of damaged, ROS-producing mitochondria is implicated in inflammasome activation, this evidence remained indirect. However, this finding suggests that MAMs may become locally enriched in ROS or ROS-derived signaling molecules and recruit ROS-sensing proteins. Consistent with this hypothesis, these authors inhibited the activity of the OMM channel VDAC and demonstrated that the knockdown of VDAC1/2 selectively abrogates NLRP3 inflammasome formation. In contrast, VDAC, and thus mitochondria, are not essential for the activation of AIM2 or IPAF inflammasomes. VDAC activity is regulated by hexokinase and Bcl-2 family members; the overexpression of Bcl-2 leads to partial VDAC closure and a concomitant decrease of mitochondrial Ca^{2+} levels and ROS production. Stimulated macrophages isolated from Bcl-2-overexpressing transgenic mice exhibit decreased levels of IL-1 β compared with cells from wild-type mice (Fig. 6).

Additional observations provide support for a pivotal role of mitochondria in NLRP3 inflammasome activation. ROS production can be specifically induced in mitochondria by inhibiting key enzymes of the electron transport chain. Addition of the complex I inhibitor rotenone results in a partial loss of Ψ_m and strong ROS production, as observed for the complex III inhibitor antimycin A; indeed, both drugs lead to NLRP3 inflammasome activation (111, 190).




FIG. 6. Schematic representation of the complex ER-mitochondria connections in the coordination of inflammasome formation. In the presence of pro-inflammatory stimuli, NLRP3 translocates to MAMs with ASC and pro-caspase1, inducing caspase-1 activation and the production of IL-1 β and IL-18. Increased levels of ROS and activation of the UPR-system *via* ER-stress are essential for NLRP3 activation. MAVS is located at the outer mitochondrial membrane and mediates antiviral signaling. ASC, apoptosis-associated speck-like protein containing a caspase recruitment domain; MAVS, mitochondria-associated viral sensor; NLRP3, NOD-like receptor family 3. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars

Furthermore, antioxidants specifically targeting mitochondria appear to block inflammasome activation and inflammation in general (20, 124).

Bulua *et al.* identified mitochondrial ROS as a driver of inflammation in tumor necrosis factor receptor-associated periodic syndrome (TRAPS), which is an autoinflammatory disorder caused by missense mutations in the type-1 TNF receptor (TNFR1), and potentially in other autoinflammatory diseases (20). When mitochondrial respiration or ROS production is inhibited, inflammatory cytokine production in response to LPS is blunted. These results suggest a more general role for mitochondrial ROS in the induction of inflammatory cytokines upstream of inflammasome activation. In TRAPS, increased mitochondrial ROS result from enhanced oxidative phosphorylation; blockade of ROS production by the mitochondria provides a new therapeutic strategy for reducing the symptoms of TRAPS and other inflammatory states.

Taken together, these and other data suggest a model in which a wide range of danger signals converge to cause increased generation of mitochondrial ROS. A rise in mitochondrial oxidant production could therefore represent the common currency of all of these divergent stress signals, with subsequent activation of the NLRP3 inflammasome. For instance, although there is compelling evidence that mitochondrial oxidants regulate the inflammasome, the mechanisms through which inflammatory signals can regulate mitochondrial function and the precise mechanism by which the release of ROS can trigger NLRP3 activation have yet to be defined. The precise molecular target of mitochondrial ROS is poorly understood, and there is little insight regarding the specific role of oxidants in inflammatory circumstances.

Another NLRP3 binding partner, TXNIP, redistributes to MAMs/mitochondria in response to oxidative stress (157) or NLRP3 inflammasome activation (189). In resting cells, TXNIP interacts with TRX and is therefore unavailable for NLRP3 interaction.

Inflammasome activators, such as uric acid crystals, induce the dissociation of TXNIP from thioredoxin in an ROS-sensitive manner and allow it to bind NLRP3 and translocate to MAMs/mitochondria (189), raising the possibility that TXNIP is involved in IL-1 β production through NLRP3 under ER stress conditions. TXNIP is a critical signaling node that links ER stress and inflammation. TXNIP is induced by ER stress through the PERK and IRE1 pathways, induces IL-1 β mRNA transcription, activates IL-1 β production by the NLRP3 inflammasome, and mediates ER stress-mediated β -cell death (134) (Fig. 6).

MAMs play a central role not only in the communication between the ER and the mitochondria but also in maintaining various cellular processes such as the antiviral response.

Recently, a new role has been identified for the adaptor MAVS (mitochondrial antiviral signaling protein) as the mitochondrial anchor for NLRP3 inflammasome formation (167). MAVS is a well-known mitochondrial protein that plays a crucial role in RIG-like receptor (RLR) signaling pathways leading to type I IFN induction and NF- κ B activation (160). Specifically, MAVS contains an N-terminal CARD-like domain and a C-terminal transmembrane domain that targets the protein to the mitochondrial membrane. The mitochondrial localization of MAVS represents the first ex-

ample of a mitochondrial protein that plays a pivotal role in innate immunity. Viral RNAs are recognized in the cytosol by the helicases RIG-1 or MDA5 (melanoma differentiation-associated gene 5). The N-termini of RIG1 and MDA5 contain two CARD domains that interact with the CARD domain of the mitochondrial adaptor MAVS. After the recruitment of transactivators, MAVS induces phosphorylation of IRF3 and IRF7 and activation of NF- κ B, leading to the production of type I IFNs and proinflammatory cytokines, respectively. It has recently been proposed that during RNA infection, RIG-1 is recruited to the MAMs to bind MAVS (77). Dynamic MAMs tethering to mitochondria and peroxisomes subsequently coordinates MAVS localization to form a signaling synapse between membranes. Importantly, the hepatitis C virus NS3/4A protease, which cleaves MAVS to support persistent infection, targets this synapse for MAVS proteolysis from the MAMs, but not from the mitochondria, to ablate RIG-1 signaling of immune responses. These results identify an innate immune signaling synapse in which the MAMs serves as the central scaffold that coordinates MAVS-dependent signaling of the RIG-1 pathway between mitochondria and peroxisomes (Fig. 6). Collectively with the role of the MAMs in NLRP3 inflammasome signaling (190), these findings indicate that the MAMs plays a central role in initiating both the innate immune and the inflammatory responses to infection.

Recently, Jacobs *et al.* demonstrated that gp78, which is an E3 ubiquitin ligase active in the ER-associated degradation (ERAD) pathway and localizes to the ER-mitochondria interface, is a novel regulator of RLR signaling (83). In addition to the enteroviruses coxsackie virus B and poliovirus, the depletion of gp78 results in a robust decrease of vesicular stomatitis virus infection and a corresponding enhancement of type I IFN signaling. Mechanistically, gp78 modulates type I IFN induction, altering both the expression and signaling of MAVS. These studies indicate an unexpected role for MAM-localized gp78 E3 ubiquitin ligase in the negative regulation of MAVS signaling. These results suggest two parallel pathways by which gp78 regulates MAVS expression and signaling: One pathway requires its E3 ubiquitin ligase and ERAD activity, whereas the second pathway requires the gp78 C-terminus and occurs *via* the association between this region and the N- and C-terminal domains of MAVS.

In addition to RIG-1, several other proteins that function in the area surrounding the mitochondria fine-tune the activities and functions of MAVS. Among these proteins, STING (STimulator of INterferon Genes) is particularly interesting, as it is enriched at MAMs; this protein interacts with RIG-1 and binds MAVS to activate a TBK-1- and IRF3-dependent cascade that ultimately induces the expression of type I IFN (82).

These results suggest that MAMs delineate a signaling synapse (involving both mitochondrial and ER components) that is essential for optimal antiviral responses.

Recently, it has been demonstrated that MAVS is required for optimal NLRP3 activity, mediating recruitment of NLRP3 to mitochondria, as well as promoting the production of IL-1 β and the pathophysiological activity of the NLRP3 inflammasome (167). The recruitment, which depends on a short N-terminal sequence in NLRP3, promotes ASC "speckle" formation and the downstream biochemical

events associated with the activity of the inflammasome. Contrary to these results, which support a role for MAVS in the activation of the NLRP3 inflammasome *via* nonviral stimuli such as LPS plus nigericin or LPS plus ATP, Park *et al.* demonstrated that MAVS regulates NLRP3 activation primarily in response to stimuli that directly engage MAVS, such as infection with Sendai virus (140). Activation of MAVS signaling by Sendai virus infection promotes NLRP3-dependent caspase-1 activation, whereas knockdown of MAVS expression clearly attenuated the activation of NLRP3 inflammasomes in THP-1 and mouse macrophages (Fig. 6). These results suggest that MAVS facilitates the recruitment of NLRP3 to the mitochondria and may enhance its oligomerization and activation by bringing it in close proximity to mitochondrial ROS.

Given the important roles of the MAMs in different cellular processes, it is not surprising that numerous viral proteins target this structure. One well-characterized example is the human cytomegalovirus glycoprotein UL37 exon 1 (16) that traffics into the MAMs during permissive infection and induces alteration of the MAMs protein composition. This glycoprotein targets MAMs with two mitochondrial targeting signals (150) and is able to reduce ER Ca^{2+} contents, possibly by modulating the amount of Ca^{2+} -regulating chaperones and oxidoreductases such as BiP present on MAMs (188) or by increasing the targeting of GRP75 to the VDAC/IP3R/GRP75 ternary complex (16).

More recently, human immunodeficiency virus type 1 (HIV-1) viral protein R (Vpr) has been suggested to be present on the ER, MAMs, and OMM, possibly *via* the integration of its C-terminal transmembrane domain (80). Vpr injures OMM and causes loss of Ψ_m by post-transcriptionally reducing the expression of MFN2 and increasing mitochondrial deformation. Vpr also markedly decreases cytoplasmic levels of DRP1 and increases bulging in MAMs. Collectively, these results suggest that Vpr-mediated cellular damage may occur *via* an alternative protein transport pathway from the ER through the MAMs to the mitochondria, which is modulated by MFN2 and DRP1 (Fig. 6).

As we have outlined thus far, MAMs play a central role in the formation and regulation of the NLRP3 inflammasome. In addition to the two aforementioned nigericin and crystalline activators, several stimuli can trigger inducible activation of the NLRP3 complex, including extracellular ATP released by dying cells (117), cholesterol crystals (44), RNA and Ca^{2+} (85).

Murakami *et al.* elucidated a critical role for Ca^{2+} mobilization in the activation of the NLRP3 inflammasome by multiple stimuli (123). These authors demonstrated that blocking Ca^{2+} mobilization inhibits the assembly and activation of the NLRP3 inflammasome complex and that during ATP stimulation, Ca^{2+} signaling is pivotal in promoting mitochondrial damage. C/EBP homologous protein, which is a transcription factor that can modulate Ca^{2+} release from the ER, amplifies NLRP3 inflammasome activation, thus linking ER stress to the activation of the NLRP3 inflammasome. Ca^{2+} signaling is not sufficient for NLRP3 inflammasome activation, as indicated by the inability of the Ca^{2+} ionophore ionomycin to induce IL-1 β production; however, only stimuli that mobilize Ca^{2+} in a manner leading to mitochondrial damage would activate the NLRP3 inflammasome.

Lee *et al.* recently confirmed and updated the role of Ca^{2+} in inflammation; these authors found that the murine Ca^{2+} -

sensing receptor (CaSR) strongly activates the NLRP3 inflammasome, which is mediated by increased intracellular Ca^{2+} and decreased cellular cyclic AMP (105). CaSR activates the NLRP3 inflammasome through phospholipase C, which catalyzes inositol-1,4,5-trisphosphate production, thereby inducing release of Ca^{2+} from ER stores; the increased cytoplasmic Ca^{2+} promotes the assembly of inflammasome components. Moreover, G-protein-coupled receptors can activate the inflammasome, indicating that increased extracellular Ca^{2+} functions as a specific amplifier of inflammation (155). Activation is mediated by signaling through the Ca^{2+} -signaling receptor and GPRC6A *via* the phosphatidylinositol/ Ca^{2+} pathway.

Recently, Nakahira *et al.* demonstrated that treatment with LPS and ATP release mtDNA into the cytosolic compartment and that this requires activation of the NLRP3 inflammasome, directly contributing to downstream activation of caspase-1 (124) (Fig. 6).

Recently, Oelze *et al.* demonstrated that glutathione peroxidase-1 (GPx-1) ablation in aging animals has a substantial impact on the burden of oxidative stress and injury (130). They observed that the age-dependent increase in the infiltration of cardiovascular tissue with leukocytes is more pronounced in GPx-1-deficient mice, suggesting that GPx-1 deficiency may lead to an inflammatory phenotype of the vasculature, which is a condition that has been reported to contribute to increased oxidative stress and vascular/endothelial dysfunction (178).

These findings give rise to further speculations. For example, MAMs may be a key player in aging-related, low-grade inflammation, as aging is associated with increased mitochondrial ROS formation, and MAMs contain several redox-sensitive components of the NLRP3 inflammasome.

Collectively, these findings highlight the central role of MAMs in the coordination of inflammasome formation and antiviral immunity. Future studies are necessary to fully explore the biological significance of MAMs during these processes.

MAMs Deregulation in the Pathogenesis of Neurological Disorders

Alterations in mitochondrial and ER homeostasis and the link between MAMs homeostasis and cellular derangement are common features of several neuronal diseases in which genetic models and environmental factors permitted the identification of common traits in the pathogenic routes. Indeed, with the increasing amount of information on MAMs, considerable evidence indicates that MAMs play an important role in neuronal disease. Analysis of the proteins present at MAMs through a mass spectrometry-based proteomic characterization from mouse brain tissue was conducted in association with a quantitative validation method to distinguish true MAMs proteins from contaminating proteins. Poston *et al.* (147) identified several proteins related to different neuronal-based diseases, such as movement disorders (chorea and Parkinson's disease), genetic disorders (Huntington's disease), and neurodegenerative diseases (schizophrenia, dementia, and seizures). This evidence suggests that several neuronal disorders share an alteration of MAMs homeostasis, where two crucial organelles (the mitochondrion and the ER) for neuronal cells are in close contact with a

sustained cross-talk. We can therefore speculate that neuronal diseases are MAMs-related disorders.

Here, we briefly discuss alterations in MAMs functions that appear to be important in the pathogenesis of selected human neuronal pathologies. We will summarize only the possible link between MAMs and these pathologies, referring readers to the literature for details of the individual pathologies.

Ca^{2+} homeostasis represents a link between MAMs and neuronal diseases, particularly through the alterations in Ca^{2+} cross-talk between the ER and mitochondria at MAMs, which are a hotspot of Ca^{2+} signaling domains (142).

Various experimental observations have suggested that an alteration of intracellular Ca^{2+} homeostasis contributes to the development of FAD and, more generally, to the pathogenesis of AD. Indeed, mutations in PS have been shown to alter ER Ca^{2+} release, affecting the mitochondria in a number of cell models (69, 106, 108). Interestingly, PSs are highly enriched in MAMs (158).

Different, and, in some cases, contrasting, hypotheses have been proposed. Some authors argue that ER Ca^{2+} overload is because wild-type PSs, but not the FAD mutants, can form Ca^{2+} -permeable leak channels in the ER (173), thus providing a clear case for enhanced Ca^{2+} release in their pathological model. Subsequent studies from other groups have presented results that are not entirely consistent with this hypothesis (26, 49, 57, 92, 187). This experimental discrepancy has been explained in part by the observation that PS1 and PS2 play distinct roles: PS2, but not PS1, modulates the ER-mitochondria tethering by increasing the number and/or the extent of their contact sites and, in turn, the Ca^{2+} cross-talk between the ER and mitochondria. FAD-linked PS2 mutations lead to a larger increase in ER-mitochondria interactions and, consequently, to an altered (and deleterious) Ca^{2+} transfer from the ER to mitochondria (Fig. 7).

There is currently a consensus that an increased association between MAMs and mitochondria is linked to the pathogenesis of AD; indeed, in different models of AD, the area of close apposition between mitochondria and the ER

(i.e., the MAMs) and the processes that occur at MAMs are enhanced compared with wild-type cells (158). Consistent with these observations, γ -secretase and PS activity is lower in MFN2-deficient cells that contain very few MAMs than in wild-type cells (5). Interestingly, this occurs not only in FAD but also in sporadic AD. However, these results were obtained in only fibroblasts from patients, not in neurons.

Several studies support the theory that the neuronal demise is potentiated by vascular alterations in the early stages of the disease. Recently, it has been demonstrated that amyloid- β ($\text{A}\beta$) induces ER stress in brain endothelial cells and triggers a mitochondria-mediated apoptotic cell death pathway involving ER-to-mitochondria Ca^{2+} transfer, decrease of Ψ_m , and release of proapoptotic factors (50), thus suggesting the participation of MAMs in this pathogenetic route (Fig. 7).


Regardless of how the synergistic " Ca^{2+} hit" occurs, mitochondrial dysfunction appears to be an obligatory downstream step in the pathogenesis of AD. Electron microscopy analysis of mitochondria in various regions of AD-affected brains revealed significant morphological organelle alterations (6).

$\text{A}\beta$ peptides increased neuronal ROS production, activated the mitochondrial fission proteins DRP1 and FIS1, and caused mitochondrial fragmentation (7).

Moreover, $\text{A}\beta$ peptides modulate the mPTP (Fig. 7). $\text{A}\beta$ peptides have been shown to inhibit mitochondrial respiration (34) and facilitate the opening of the mPTP. However, knockout mouse models for cyclophilin D, which is an essential component of mPTP activity, exhibit improved cognitive abilities and a minor $\text{A}\beta$ -mediated reduction of long-term potentiation (43).

MAMs also play a key role in Parkinson's disease (PD) (Fig. 7). This hypothesis is supported by the identification of a cohort of proteins involved in the familial forms of PD that appear to share intracellular localization of MAMs and possibly indicate a signaling role for Ca^{2+} . Specifically, mutations were reported in genes encoding for α -synuclein, DJ-1, PINK1, and parkin.

FIG. 7. Involvement of the ER-mitochondria interface in the pathogenesis of neuronal disorders such as Alzheimer's disease (AD) and Parkinson's disease (PD). Alterations of Ca^{2+} homeostasis represent a link between MAMs and neuronal diseases, particularly regarding the alterations of Ca^{2+} crosstalk between the ER and mitochondria at the MAMs. $\text{A}\beta$ oligomers affect mitochondrial functionality by enhancing ER Ca^{2+} and facilitate the opening of the PTP. α -synuclein positively affects Ca^{2+} transfer from the ER to the mitochondria. $\text{A}\beta$, amyloid-beta; cyt C, cytochrome C; PSs, presenilins. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars



α -Synuclein is localized to mitochondria-associated ER membranes (68). The different levels of α -synuclein oligomerization have been linked to cell death. In particular, a heterogeneous mixture of small oligomers of α -synuclein can lead to Ca^{2+} dysregulation to the point of mPTP activation and commitment to neuronal cell death (36). Studies have shown that α -synuclein positively affects Ca^{2+} transfer from the ER to the mitochondria (22, 89, 119, 161). This effect is correlated with an increase in the number of MAMs through the C-terminal of the α -synuclein domain (22). Moreover, it has been proposed that the accumulation of α -synuclein in the cells causes the redistribution of α -synuclein to localized foci and reduces the ability of mitochondria to accumulate Ca^{2+} , resulting in augmented autophagy with the risk of altering normal mitochondrial homeostasis (22).

DJ-1 is largely cytoplasmic, except for a pool localized in mitochondria, most likely within the IMS and in the matrix; whereas little, if any, DJ-1 is associated with the outer and inner mitochondrial membranes (41). Numerous studies support the role of DJ-1 in mitigating oxidative stress and in the maintenance of mitochondrial homeostasis (41). A recent study proposed that DJ-1 modulates ER-mitochondrial Ca^{2+} cross-talk, favoring the tethering between the two organelles and, thus, MAMs formation (135).

Parkin is a ubiquitin-protein ligase that localizes to the mitochondrial matrix, where it enhances mitochondrial gene transcription and biogenesis in proliferating cells (84). Parkin has been proposed to play a neuroprotective role, promoting the clearance of damaged mitochondria through the mitophagic process (127). In *Drosophila*, parkin null mutants exhibit defects in mitochondrial function and increased oxidative stress. In contrast, overexpression of parkin in cultured cells prevents mitochondrial swelling and stress-induced apoptosis (84). Parkin overexpression stimulates MAMs formation and, in turn, physically and functionally enhances ER-mitochondria coupling, thus

favoring Ca^{2+} transfer from the ER to the mitochondria and energy metabolism (23).

As previously described, parkin associates with PINK1 in mitochondrial quality control pathways, promoting the selective degradation of damaged mitochondria through mitophagy (175, 185).

PINK1 is unambiguously localized to mitochondrial membranes, and PINK1 overexpression protects cells from mitochondrial depolarization and apoptosis (183). Loss-of-function mutations are responsible for dopaminergic neuronal degeneration in *Drosophila* (139). Overexpression of parkin has been shown to rescue the mitochondrial dysfunction caused by PINK1 deficiency (184).

Finally, a mutant form of PINK1 has been shown to exacerbate mitochondrial alterations (disturbing mitochondrial Ca^{2+} fluxes) promoted by an α -synuclein mutant, thus suggesting cooperative roles for these two proteins (118).

GM1-gangliosidosis represents an additional example of a MAMs-related pathology. This neurodegenerative disease is characterized by GM1-ganglioside (GM1) accumulation within MAMs in brain tissue (156).

In addition, series of independent observations highlighted a possible role for MAMs in Huntington's disease. Mutant, but not wild-type huntingtin (Htt; the protein that, when mutated, is responsible for the disease), localizes to the mitochondrial membranes in neurons (136). Htt forms a ternary complex with Htt-associated protein-1A (HAP-1A) and IP3R (a protein enriched at MAMs). In this complex, mutant Htt, but not wild-type Htt, facilitates Ca^{2+} release from the ER and renders neurons more sensitive to Ca^{2+} -mediated cellular dysfunction *via* mitochondrial Ca^{2+} overload (9) and mPTP opening (Fig. 8) (27).

MAMs involvement appears to also be important in amyotrophic lateral sclerosis (ALS). Vesicle-associated membrane protein-associated protein B (VAPB) has been proposed to play a role in the pathogenesis of this neurological disease,





FIG. 8. Schematic representation of the ER-mitochondria interface in Huntington's disease (HD), amyotrophic lateral sclerosis (ALS), and Wolfram syndrome. In HD, mutant HTT increases ER Ca^{2+} release by acting on InsP_3R . Mutant VAPB enhances ER-mitochondria tethering in ALS; Sig1R has also been proposed to be involved in the pathogenesis of ALS. Miner1, a protein that, when mutated, causes Wolfram syndrome, is enriched at MAMs. HAP-1A, huntingtin-associated protein 1-A; mutHTT, mutant huntingtin; PTPIP51, protein tyrosine phosphatase-interacting protein 51; VAPB, vesicle-associated membrane protein-associated protein B. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars

FIG. 9. An illustrated overview of the main functions attributed to MAMs. Correct homeostasis of MAMs is a key determinant of several cellular processes. For example, the activities of several MAM proteins have been found to be corrupted during the pathogenesis of a number of human diseases. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars



and recent studies localize this protein at MAMs, where it interacts with protein tyrosine phosphatase interacting protein 51 (PTPIP51), which is an outer mitochondrial membrane protein, and regulates ER-mitochondrial Ca²⁺ cross-talk. The VAPBP mutant associated with ALS has been demonstrated to alter its binding to PTPIP51 and exacerbate ER-mitochondrial Ca²⁺ transfer (39) (Fig. 8).

Sig1R, which is another important protein that is particularly enriched at MAMs and is involved in Ca²⁺ homeostasis, has been proposed to be involved in the pathogenesis of ALS (148) (Fig. 8).

Interestingly, in several studies, mitochondrial dysfunction has been reported to be frequently associated with demyelination, whereas proper mitochondrial function is required for correct oligodendrocyte differentiation and myelination. Bonora and colleagues reported that correct mitochondrial Ca²⁺ signaling and, thus, MAMs activity are impaired in conditions mimicking the proinflammatory environment to which the oligodendrocytes are exposed in multiple sclerosis patients. These abnormalities result in inefficient oligodendrocyte differentiation (13).

Finally, MAMs are also emerging as critical intracellular domains in the pathogenesis of the incurable disease Wolfram syndrome. This concept is demonstrated by the observation that Miner1, which is a protein that, if mutated, causes Wolfram syndrome, is enriched at MAMs, and is involved in ER stress and Ca²⁺ signaling around MAMs, as previously described for other proteins involved in mitochondrial structure and physiology (180).

These results highlight MAMs involvement in the pathogenesis of several neurological disorders. Studies on this topic are ongoing in several groups, including ours, and we predict that in the future, new findings will consolidate this concept for the disorders presented here and for numerous others.

Concluding Remarks

Overall, the picture emerging from the study of MAMs appears to be extremely complex (Fig. 9), with several un-

certainities to be resolved. Nevertheless, the common role of MAMs in important physiopathological pathways is clear, providing a leading theme for future studies. However, whether alterations in MAMs represent a response to the disease pathogenesis or directly contribute to the disease has not yet been unequivocally established. In any case, MAMs represent a promising pharmacological target for several important human diseases. Finally, the following important questions remain open: (i) What are the entire proteomes of the MAMs in the different cell and tissue types under normal conditions and under stress or pathological conditions? (ii) Are the MAMs identical within the cell, or is any cell endowed with several different types of MAMs with respect to protein composition? (iii) What are the precise molecular mechanisms of MAMs formation? and (iv) How are the proteins and lipids targeted to MAMs?

Acknowledgments

This study was supported by the Italian Association for Cancer Research (AIRC) and local funding from the University of Ferrara to Paolo Pinton and Carlotta Giorgi; and from Telethon (GGP11139B), the Italian Cystic Fibrosis Foundation (Grant 19/2014), the Italian Ministry of Health, the Italian Ministry of Education, and the University and Research (COFIN, FIRB, and Futuro in Ricerca) to Paolo Pinton. Mariusz R. Wieckowski and Jerzy Duszynski were supported by the Polish National Science Center (UMO-2011/11/M/NZ3/02128), the Polish Ministry of Science and Higher Education grant W100/HFSC/2011 and Grant HFSP RGP0027/2011. Simone Patergnani was supported by a research fellowship from FISM (Fondazione Italiana Sclerosi Multipla), cod. 2012/B/11.

References

1. Alberts B. The cell as a collection of protein machines: preparing the next generation of molecular biologists. *Cell* 92: 291–294, 1998.

2. Anelli T, Bergamelli L, Margittai E, Rimessi A, Fagioli C, Malgaroli A, Pinton P, Ripamonti M, Rizzuto R, and Sitia R. Ero1alpha regulates Ca(2+) fluxes at the endoplasmic reticulum-mitochondria interface (MAM). *Antioxid Redox Signal* 16: 1077–1087, 2012.
3. Appenzeller-Herzog C, Riemer J, Zito E, Chin KT, Ron D, Spiess M, and Ellgaard L. Disulphide production by Ero1alpha-PDI relay is rapid and effectively regulated. *EMBO J* 29: 3318–3329, 2010.
4. Area-Gomez E, de Groof AJ, Boldogh I, Bird TD, Gibson GE, Koehler CM, Yu WH, Duff KE, Yaffe MP, Pon LA, and Schon EA. Presenilins are enriched in endoplasmic reticulum membranes associated with mitochondria. *Am J Pathol* 175: 1810–1816, 2009.
5. Area-Gomez E, Del Carmen Lara Castillo M, Tambini MD, Guardia-Laguarta C, de Groof AJ, Madra M, Ikenouchi J, Umeda M, Bird TD, Sturley SL, and Schon EA. Upregulated function of mitochondria-associated ER membranes in Alzheimer disease. *EMBO J* 31: 4106–4123, 2012.
6. Baloyannis SJ. Mitochondria are related to synaptic pathology in Alzheimer's disease. *Int J Alzheimers Dis* 2011: 305395, 2011.
7. Barsoum MJ, Yuan H, Gerencser AA, Liot G, Kushnareva Y, Graber S, Kovacs I, Lee WD, Waggoner J, Cui J, White AD, Bossy B, Martinou JC, Youle RJ, Lipton SA, Ellisman MH, Perkins GA, and Bossy-Wetzel E. Nitric oxide-induced mitochondrial fission is regulated by dynamin-related GTPases in neurons. *EMBO J* 25: 3900–3911, 2006.
8. Betz C, Stracka D, Prescianotto-Baschong C, Frieden M, Demaurex N, and Hall MN. Feature article: mTOR complex 2-Akt signaling at mitochondria-associated endoplasmic reticulum membranes (MAM) regulates mitochondrial physiology. *Proc Natl Acad Sci U S A* 110: 12526–12534, 2013.
9. Bezprozvanny I and Hayden MR. Deranged neuronal calcium signaling and Huntington disease. *Biochem Biophys Res Commun* 322: 1310–1317, 2004.
10. Bononi A, Agnoletto C, De Marchi E, Marchi S, Patergnani S, Bonora M, Giorgi C, Missiroli S, Poletti F, Rimessi A, and Pinton P. Protein kinases and phosphatases in the control of cell fate. *Enzyme Res* 2011: 329098, 2011.
11. Bononi A, Bonora M, Marchi S, Missiroli S, Poletti F, Giorgi C, Pandolfi PP, and Pinton P. Identification of PTEN at the ER and MAMs and its regulation of Ca(2+) signaling and apoptosis in a protein phosphatase-dependent manner. *Cell Death Differ* 20: 1631–1643, 2013.
12. Bonora M, Bononi A, De Marchi E, Giorgi C, Lebedzinska M, Marchi S, Patergnani S, Rimessi A, Suski JM, Wojtala A, Wieckowski MR, Kroemer G, Galluzzi L, and Pinton P. Role of the c subunit of the FO ATP synthase in mitochondrial permeability transition. *Cell Cycle* 12: 674–683, 2013.
13. Bonora M, De Marchi E, Patergnani S, Suski JM, Celsi F, Bononi A, Giorgi C, Marchi S, Rimessi A, Duszynski J, Pozzan T, Wieckowski MR, and Pinton P. Tumor necrosis factor-alpha impairs oligodendroglial differentiation through a mitochondria-dependent process. *Cell Death Differ* 21: 1198–1208, 2014.
14. Bonora M, Patergnani S, Rimessi A, De Marchi E, Suski JM, Bononi A, Giorgi C, Marchi S, Missiroli S, Poletti F, Wieckowski MR, and Pinton P. ATP synthesis and storage. *Purinergic Signal* 8: 343–357, 2012.
15. Bonora M, Wieckowski MR, Chinopoulos C, Kepp O, Kroemer G, Galluzzi L, and Pinton P. Molecular mechanisms of cell death: central implication of ATP synthase in mitochondrial permeability transition. *Oncogene* 0, 2014.
16. Bozidis P, Williamson CD, Wong DS, and Colberg-Poley AM. Trafficking of UL37 proteins into mitochondrion-associated membranes during permissive human cytomegalovirus infection. *J Virol* 84: 7898–7903, 2010.
17. Bravo R, Vicencio JM, Parra V, Troncoso R, Munoz JP, Bui M, Quiroga C, Rodriguez AE, Verdejo HE, Ferreira J, Iglewski M, Chiong M, Simmen T, Zorzano A, Hill JA, Rothermel BA, Szabadkai G, and Lavandero S. Increased ER-mitochondrial coupling promotes mitochondrial respiration and bioenergetics during early phases of ER stress. *J Cell Sci* 124: 2143–2152, 2011.
18. Brooks C, Wei Q, Feng L, Dong G, Tao Y, Mei L, Xie ZJ, and Dong Z. Bak regulates mitochondrial morphology and pathology during apoptosis by interacting with mitofusins. *Proc Natl Acad Sci U S A* 104: 11649–11654, 2007.
19. Brown GC and Borutaite V. There is no evidence that mitochondria are the main source of reactive oxygen species in mammalian cells. *Mitochondrion* 12: 1–4, 2012.
20. Bulua AC, Simon A, Maddipati R, Pelletier M, Park H, Kim KY, Sack MN, Kastner DL, and Siegel RM. Mitochondrial reactive oxygen species promote production of proinflammatory cytokines and are elevated in TNFR1-associated periodic syndrome (TRAPS). *J Exp Med* 208: 519–533, 2011.
21. Cabibbo A, Pagani M, Fabbri M, Rocchi M, Farmery MR, Bulleid NJ, and Sitia R. ERO1-L, a human protein that favors disulfide bond formation in the endoplasmic reticulum. *J Biol Chem* 275: 4827–4833, 2000.
22. Cali T, Ottolini D, Negro A, and Brini M. alpha-Synuclein controls mitochondrial calcium homeostasis by enhancing endoplasmic reticulum-mitochondria interactions. *J Biol Chem* 287: 17914–17929, 2012.
23. Cali T, Ottolini D, Negro A, and Brini M. Enhanced parkin levels favor ER-mitochondria crosstalk and guarantee Ca(2+) transfer to sustain cell bioenergetics. *Biochim Biophys Acta* 1832: 495–508, 2013.
24. Camici GG, Schiavoni M, Francia P, Bachschmid M, Martin-Padura I, Hersberger M, Tanner FC, Pelicci P, Volpe M, Anversa P, Luscher TF, and Cosentino F. Genetic deletion of p66(Shc) adaptor protein prevents hyperglycemia-induced endothelial dysfunction and oxidative stress. *Proc Natl Acad Sci U S A* 104: 5217–5222, 2007.
25. Chami M, Oules B, Szabadkai G, Tacine R, Rizzuto R, and Paterlini-Brechot P. Role of SERCA1 truncated isoform in the proapoptotic calcium transfer from ER to mitochondria during ER stress. *Mol Cell* 32: 641–651, 2008.
26. Cheung KH, Shineman D, Muller M, Cardenas C, Mei L, Yang J, Tomita T, Iwatsubo T, Lee VM, and Foscett JK. Mechanism of Ca²⁺ disruption in Alzheimer's disease by presenilin regulation of InsP3 receptor channel gating. *Neuron* 58: 871–883, 2008.
27. Choo YS, Johnson GV, MacDonald M, Detloff PJ, and Lesort M. Mutant huntingtin directly increases susceptibility of mitochondria to the calcium-induced permeability transition and cytochrome c release. *Hum Mol Genet* 13: 1407–1420, 2004.
28. Cipolat S, Martins de Brito O, Dal Zilio B, and Scorrano L. OPA1 requires mitofusin 1 to promote mitochondrial fusion. *Proc Natl Acad Sci U S A* 101: 15927–15932, 2004.
29. Colombi M, Molle KD, Benjamin D, Rattenbacher-Kiser K, Schaefer C, Betz C, Thiemeyer A, Regenass U, Hall

- MN, and Moroni C. Genome-wide shRNA screen reveals increased mitochondrial dependence upon mTORC2 addiction. *Oncogene* 30: 1551–1565, 2011.
30. Cook KL, Soto-Pantoja DR, Abu-Asab M, Clarke PA, Roberts DD, and Clarke R. Mitochondria directly donate their membrane to form autophagosomes during a novel mechanism of parkin-associated mitophagy. *Cell Biosci* 4: 16, 2014.
 31. Copeland DE and Dalton AJ. An association between mitochondria and the endoplasmic reticulum in cells of the pseudobranch gland of a teleost. *J Biophys Biochem Cytol* 5: 393–396, 1959.
 32. Cosentino F, Francia P, Camici GG, Pelicci PG, Luscher TF, and Volpe M. Final common molecular pathways of aging and cardiovascular disease: role of the p66Shc protein. *Arterioscler Thromb Vasc Biol* 28: 622–628, 2008.
 33. Cosentino K and Garcia-Saez AJ. Mitochondrial alterations in apoptosis. *Chem Phys Lipids* 181: 62–75, 2014.
 34. Crouch PJ, Blake R, Duce JA, Ciccotosto GD, Li QX, Barnham KJ, Curtain CC, Cherny RA, Cappai R, Dyrks T, Masters CL, and Trounce IA. Copper-dependent inhibition of human cytochrome c oxidase by a dimeric conformer of amyloid-beta1–42. *J Neurosci* 25: 672–679, 2005.
 35. Dallaporta B, Marchetti P, de Pablo MA, Maise C, Duc HT, Metivier D, Zamzami N, Geuskens M, and Kroemer G. Plasma membrane potential in thymocyte apoptosis. *J Immunol* 162: 6534–6542, 1999.
 36. Danzer KM, Haasen D, Karow AR, Moussaud S, Habeck M, Giese A, Kretschmar H, Hengerer B, and Kostka M. Different species of alpha-synuclein oligomers induce calcium influx and seeding. *J Neurosci* 27: 9220–9232, 2007.
 37. De Marchi E, Baldassari F, Bononi A, Wieckowski MR, and Pinton P. Oxidative stress in cardiovascular diseases and obesity: role of p66Shc and protein kinase C. *Oxid Med Cell Longev* 2013: 564961, 2013.
 38. De Stefani D, Bononi A, Romagnoli A, Messina A, De Pinto V, Pinton P, and Rizzuto R. VDAC1 selectively transfers apoptotic Ca²⁺ signals to mitochondria. *Cell Death Differ* 19: 267–273, 2012.
 39. De Vos KJ, Morotz GM, Stoica R, Tudor EL, Lau KF, Ackerley S, Warley A, Shaw CE, and Miller CC. VAPB interacts with the mitochondrial protein PTPIP51 to regulate calcium homeostasis. *Hum Mol Genet* 21: 1299–1311, 2012.
 40. Decuyper JP, Monaco G, Bultynck G, Missiaen L, De Smedt H, and Parys JB. The IP(3) receptor-mitochondria connection in apoptosis and autophagy. *Biochim Biophys Acta* 1813: 1003–1013, 2011.
 41. Dodson MW and Guo M. Pink1, Parkin, DJ-1 and mitochondrial dysfunction in Parkinson's disease. *Curr Opin Neurobiol* 17: 331–337, 2007.
 42. Drose S, Brandt U, and Wittig I. Mitochondrial respiratory chain complexes as sources and targets of thiol-based redox-regulation. *Biochim Biophys Acta* 1844: 1344–1354, 2014.
 43. Du H, Guo L, Fang F, Chen D, Sosunov AA, McKhann GM, Yan Y, Wang C, Zhang H, Molkenstein JD, Gunn-Moore FJ, Vonsattel JP, Arancio O, Chen JX, and Yan SD. Cyclophilin D deficiency attenuates mitochondrial and neuronal perturbation and ameliorates learning and memory in Alzheimer's disease. *Nat Med* 14: 1097–1105, 2008.
 44. Duewell P, Kono H, Rayner KJ, Sirois CM, Vladimer G, Bauernfeind FG, Abela GS, Franchi L, Nunez G, Schnurr M, Espevik T, Lien E, Fitzgerald KA, Rock KL, Moore KJ, Wright SD, Hornung V, and Latz E. NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature* 464: 1357–1361, 2010.
 45. Ellgaard L and Ruddock LW. The human protein disulphide isomerase family: substrate interactions and functional properties. *EMBO Rep* 6: 28–32, 2005.
 46. Ellis RJ. Macromolecular crowding: obvious but underappreciated. *Trends Biochem Sci* 26: 597–604, 2001.
 47. Enyedi B, Varnai P, and Geiszt M. Redox state of the endoplasmic reticulum is controlled by Ero1L-alpha and intraluminal calcium. *Antioxid Redox Signal* 13: 721–729, 2010.
 48. Eskes R, Antonsson B, Osen-Sand A, Montessuit S, Richter C, Sadoul R, Mazzei G, Nichols A, and Martinou JC. Bax-induced cytochrome C release from mitochondria is independent of the permeability transition pore but highly dependent on Mg²⁺ ions. *J Cell Biol* 143: 217–224, 1998.
 49. Fedrizzi L, Lim D, Carafoli E, and Brini M. Interplay of the Ca²⁺-binding protein DREAM with presenilin in neuronal Ca²⁺ signaling. *J Biol Chem* 283: 27494–27503, 2008.
 50. Fonseca AC, Moreira PI, Oliveira CR, Cardoso SM, Pinton P, and Pereira CF. Amyloid-beta disrupts calcium and redox homeostasis in brain endothelial cells. *Mol Neurobiol*, 2014 [Epub ahead of print]; DOI: 10.1007/s12035-014-8740-7
 51. Frank S, Gaume B, Bergmann-Leitner ES, Leitner WW, Robert EG, Catez F, Smith CL, and Youle RJ. The role of dynamin-related protein 1, a mediator of mitochondrial fission, in apoptosis. *Dev Cell* 1: 515–525, 2001.
 52. Fujimoto M, Hayashi T, and Su TP. The role of cholesterol in the association of endoplasmic reticulum membranes with mitochondria. *Biochem Biophys Res Commun* 417: 635–639, 2012.
 53. Garcia Fernandez M, Troiano L, Moretti L, Nasi M, Pinti M, Salvioli S, Dobrucki J, and Cossarizza A. Early changes in intramitochondrial cardiolipin distribution during apoptosis. *Cell Growth Differ* 13: 449–455, 2002.
 54. Gegg ME, Cooper JM, Chau KY, Rojo M, Schapira AH, and Taanman JW. Mitofusin 1 and mitofusin 2 are ubiquitinated in a PINK1/parkin-dependent manner upon induction of mitophagy. *Hum Mol Genet* 19: 4861–4870, 2010.
 55. Geisler S, Holmstrom KM, Skujat D, Fiesel FC, Rothfuss OC, Kahle PJ, and Springer W. PINK1/Parkin-mediated mitophagy is dependent on VDAC1 and p62/SQSTM1. *Nat Cell Biol* 12: 119–131, 2010.
 56. Gertz M, Fischer F, Wolters D, and Steegborn C. Activation of the lifespan regulator p66Shc through reversible disulfide bond formation. *Proc Natl Acad Sci U S A* 105: 5705–5709, 2008.
 57. Giacomello M, Barbiero L, Zatti G, Squitti R, Binetti G, Pozzan T, Fasolato C, Ghidoni R, and Pizzo P. Reduction of Ca²⁺ stores and capacitative Ca²⁺ entry is associated with the familial Alzheimer's disease presenilin-2 T122R mutation and anticipates the onset of dementia. *Neurobiol Dis* 18: 638–648, 2005.
 58. Gilady SY, Bui M, Lynes EM, Benson MD, Watts R, Vance JE, and Simmen T. Ero1alpha requires oxidizing and normoxic conditions to localize to the mitochondria-associated membrane (MAM). *Cell Stress Chaperones* 15: 619–629, 2010.
 59. Giorgi C, Agnoletto C, Baldini C, Bononi A, Bonora M, Marchi S, Missiroli S, Patergnani S, Poletti F, Rimessi A, Zavan B, and Pinton P. Redox control of protein kinase C: cell- and disease-specific aspects. *Antioxid Redox Signal* 13: 1051–1085, 2010.

60. Giorgi C, Baldassari F, Bononi A, Bonora M, De Marchi E, Marchi S, Missiroli S, Patergnani S, Rimessi A, Suski JM, Wieckowski MR, and Pinton P. Mitochondrial Ca(2+) and apoptosis. *Cell Calcium* 52: 36–43, 2012.
61. Giorgi C, De Stefani D, Bononi A, Rizzuto R, and Pinton P. Structural and functional link between the mitochondrial network and the endoplasmic reticulum. *Int J Biochem Cell Biol* 41: 1817–1827, 2009.
62. Giorgi C, Ito K, Lin HK, Santangelo C, Wieckowski MR, Lebedzinska M, Bononi A, Bonora M, Duszynski J, Bernardi R, Rizzuto R, Tacchetti C, Pinton P, and Pandolfi PP. PML regulates apoptosis at endoplasmic reticulum by modulating calcium release. *Science* 330: 1247–1251, 2010.
63. Giorgi C, Bonora M, Sorrentino G, Missiroli S, Poletti F, Suski JM, Galindo Ramirez F, Rizzuto R, Di Virgilio F, Zito E, Pandolfi PP, Wieckowski MR, Mammano F, Del Sal G, Pinton P. p53 at the endoplasmic reticulum regulates apoptosis in a Ca²⁺-dependent manner. *Proc Natl Acad Sci U S A* 112: 1779–1784, 2015.
64. Giorgio M, Migliaccio E, Orsini F, Paolucci D, Moroni M, Contursi C, Pelliccia G, Luzi L, Minucci S, Marcaccio M, Pinton P, Rizzuto R, Bernardi P, Paolucci F, and Pelicci PG. Electron transfer between cytochrome c and p66Shc generates reactive oxygen species that trigger mitochondrial apoptosis. *Cell* 122: 221–233, 2005.
65. Graiani G, Lagrasta C, Migliaccio E, Spillmann F, Meloni M, Madeddu P, Quaini F, Padura IM, Lanfrancone L, Pelicci P, and Emanuelli C. Genetic deletion of the p66Shc adaptor protein protects from angiotensin II-induced myocardial damage. *Hypertension* 46: 433–440, 2005.
66. Green DR and Kroemer G. The pathophysiology of mitochondrial cell death. *Science* 305: 626–629, 2004.
67. Grisham MB. Methods to detect hydrogen peroxide in living cells: Possibilities and pitfalls. *Comp Biochem Physiol A Mol Integr Physiol* 165: 429–438, 2013.
68. Guardia-Laguarta C, Area-Gomez E, Rub C, Liu Y, Magrane J, Becker D, Voos W, Schon EA, and Przedborski S. alpha-Synuclein is localized to mitochondria-associated ER membranes. *J Neurosci* 34: 249–259, 2014.
69. Guo Q, Furukawa K, Sopher BL, Pham DG, Xie J, Robinson N, Martin GM, and Mattson MP. Alzheimer's PS-1 mutation perturbs calcium homeostasis and sensitizes PC12 cells to death induced by amyloid beta-peptide. *Neuroreport* 8: 379–383, 1996.
70. Hailey DW, Rambold AS, Satpute-Krishnan P, Mitra K, Sougrat R, Kim PK, and Lippincott-Schwartz J. Mitochondria supply membranes for autophagosome biogenesis during starvation. *Cell* 141: 656–667, 2010.
71. Halliwell B and Whiteman M. Measuring reactive species and oxidative damage *in vivo* and in cell culture: how should you do it and what do the results mean? *Br J Pharmacol* 142: 231–255, 2004.
72. Hamasaki M, Furuta N, Matsuda A, Nezu A, Yamamoto A, Fujita N, Oomori H, Noda T, Haraguchi T, Hiraoka Y, Amano A, and Yoshimori T. Autophagosomes form at ER-mitochondria contact sites. *Nature* 495: 389–393, 2013.
73. Harris MH and Thompson CB. The role of the Bcl-2 family in the regulation of outer mitochondrial membrane permeability. *Cell Death Differ* 7: 1182–1191, 2000.
74. Hatahet F and Ruddock LW. Protein disulfide isomerase: a critical evaluation of its function in disulfide bond formation. *Antioxid Redox Signal* 11: 2807–2850, 2009.
75. Hattori N, Saiki S, and Imai Y. Regulation by mitophagy. *Int J Biochem Cell Biol* 53: 147–150, 2014.
76. Higo T, Hattori M, Nakamura T, Natsume T, Michikawa T, and Mikoshiba K. Subtype-specific and ER lumenal environment-dependent regulation of inositol 1,4,5-trisphosphate receptor type 1 by ERp44. *Cell* 120: 85–98, 2005.
77. Horner SM, Liu HM, Park HS, Briley J, and Gale M, Jr. Mitochondrial-associated endoplasmic reticulum membranes (MAM) form innate immune synapses and are targeted by hepatitis C virus. *Proc Natl Acad Sci U S A* 108: 14590–14595, 2011.
78. Horvath SE and Daum G. Lipids of mitochondria. *Prog Lipid Res* 52: 590–614, 2013.
79. Hu Y, Wang X, Zeng L, Cai DY, Sabapathy K, Goff SP, Firpo EJ, and Li B. ERK phosphorylates p66shcA on Ser36 and subsequently regulates p27kip1 expression via the Akt-FOXO3a pathway: implication of p27kip1 in cell response to oxidative stress. *Mol Biol Cell* 16: 3705–3718, 2005.
80. Huang CY, Chiang SF, Lin TY, Chiou SH, and Chow KC. HIV-1 Vpr triggers mitochondrial destruction by impairing Mfn2-mediated ER-mitochondria interaction. *PLoS One* 7: e33657, 2012.
81. Ishihara N, Nomura M, Jofuku A, Kato H, Suzuki SO, Masuda K, Otera H, Nakanishi Y, Nonaka I, Goto Y, Taguchi N, Morinaga H, Maeda M, Takayanagi R, Yokota S, and Mihara K. Mitochondrial fission factor Drp1 is essential for embryonic development and synapse formation in mice. *Nat Cell Biol* 11: 958–966, 2009.
82. Ishikawa H and Barber GN. STING is an endoplasmic reticulum adaptor that facilitates innate immune signaling. *Nature* 455: 674–678, 2008.
83. Jacobs JL, Zhu J, Sarkar SN, and Coyne CB. Regulation of mitochondrial antiviral signaling (MAVS) expression and signaling by the mitochondria-associated endoplasmic reticulum membrane (MAM) protein Gp78. *J Biol Chem* 289: 1604–1616, 2014.
84. Jiang H, Ren Y, Zhao J, and Feng J. Parkin protects human dopaminergic neuroblastoma cells against dopamine-induced apoptosis. *Hum Mol Genet* 13: 1745–1754, 2004.
85. Jin C, Frayssinet P, Pelker R, Cwirka D, Hu B, Vignery A, Eisenbarth SC, and Flavell RA. NLRP3 inflammasome plays a critical role in the pathogenesis of hydroxyapatite-associated arthropathy. *Proc Natl Acad Sci U S A* 108: 14867–14872, 2011.
86. Kagan VE, Tyurin VA, Jiang J, Tyurina YY, Ritov VB, Amoscato AA, Osipov AN, Belikova NA, Kapralov AA, Kini V, Vlasova II, Zhao Q, Zou M, Di P, Svistunenko DA, Kurnikov IV, and Borisenko GG. Cytochrome c acts as a cardiolipin oxygenase required for release of proapoptotic factors. *Nat Chem Biol* 1: 223–232, 2005.
87. Kalwarczyk T, Ziebac N, Bielejewska A, Zaboklicka E, Koynov K, Szymanski J, Wilk A, Patkowski A, Gapinski J, Butt HJ, and Holyst R. Comparative analysis of viscosity of complex liquids and cytoplasm of mammalian cells at the nanoscale. *Nano Lett* 11: 2157–2163, 2011.
88. Kalyanaraman B, Darley-Usmar V, Davies KJ, Dennery PA, Forman HJ, Grisham MB, Mann GE, Moore K, Roberts LJ, 2nd, and Ischiropoulos H. Measuring reactive oxygen and nitrogen species with fluorescent probes: challenges and limitations. *Free Radic Biol Med* 52: 1–6, 2012.
89. Kamp F, Exner N, Lutz AK, Wender N, Hegermann J, Brunner B, Nuscher B, Bartels T, Giese A, Beyer K, Eimer S, Winklhofer KF, and Haass C. Inhibition of mitochondrial fusion by alpha-synuclein is rescued by PINK1, Parkin and DJ-1. *EMBO J* 29: 3571–3589, 2010.

90. Karbowski M, Arnoult D, Chen H, Chan DC, Smith CL, and Youle RJ. Quantitation of mitochondrial dynamics by photolabeling of individual organelles shows that mitochondrial fusion is blocked during the Bax activation phase of apoptosis. *J Cell Biol* 164: 493–499, 2004.
91. Karpinich NO, Tafani M, Rothman RJ, Russo MA, and Farber JL. The course of etoposide-induced apoptosis from damage to DNA and p53 activation to mitochondrial release of cytochrome c. *J Biol Chem* 277: 16547–16552, 2002.
92. Kasri NN, Kocks SL, Verbert L, Hebert SS, Callewaert G, Parys JB, Missiaen L, and De Smedt H. Up-regulation of inositol 1,4,5-trisphosphate receptor type 1 is responsible for a decreased endoplasmic-reticulum Ca^{2+} content in presenilin double knock-out cells. *Cell Calcium* 40: 41–51, 2006.
93. Kim CS, Jung SB, Naqvi A, Hoffman TA, DeRicco J, Yamamori T, Cole MP, Jeon BH, and Irani K. p53 impairs endothelium-dependent vasomotor function through transcriptional upregulation of p66shc. *Circ Res* 103: 1441–1450, 2008.
94. Klecker T, Bockler S, and Westermann B. Making connections: interorganelle contacts orchestrate mitochondrial behavior. *Trends Cell Biol* 24: 537–545, 2014.
95. Kliensky DJ, Abdalla FC, Abeliovich H, Abraham RT, Acevedo-Arozena A, Adeli K, Agholme L, Agnello M, Agostinis P, Aguirre-Ghiso JA, Ahn HJ, Ait-Mohamed O, Ait-Si-Ali S, Akematsu T, Akira S, Al-Younes HM, Al-Zeer MA, Albert ML, Albin RL, Alegre-Abarrategui J, Aleo MF, Alirezaei M, Almasan A, Almonte-Becerril M, Amano A, Amaravadi R, Amarnath S, Amer AO, Andrieu-Abadie N, Anantharam V, Ann DK, Anoopkumar-Dukie S, Aoki H, Apostolova N, Arancia G, Aris JP, Asanuma K, Asare NY, Ashida H, Askanas V, Askew DS, Auberger P, Baba M, Backues SK, Baehrecke EH, Bahr BA, Bai XY, Bailly Y, Baiocchi R, Baldini G, Balduini W, Ballabio A, Bamber BA, Bampton ET, Banhegyi G, Bartholomew CR, Bassham DC, Bast RC, Jr., Batoko H, Bay BH, Beau I, Bechet DM, Begley TJ, Behl C, Behrends C, Bekri S, Bellaire B, Bendall LJ, Benetti L, Berliocchi L, Bernardi H, Bernassola F, Besteiro S, Bhatia-Kissova I, Bi X, Biard-Piechaczyk M, Blum JS, Boise LH, Bonaldo P, Boone DL, Bornhauser BC, Bortoluci KR, Bossis I, Bost F, Bourquin JP, Boya P, Boyer-Guittaut M, Bozhkov PV, Brady NR, Brancolini C, Brech A, Brenman JE, Brennand A, Bresnick EH, Brest P, Bridges D, Bristol ML, Brookes PS, Brown EJ, Brumell JH, Brunetti-Pierri N, Brunk UT, Bulman DE, Bultman SJ, Bultynck G, Burbulla LF, Bursch W, Butchar JP, Buzgariu W, Bydowski SP, Cadwell K, Cahova M, Cai D, Cai J, Cai Q, Calabretta B, Calvo-Garrido J, Camougrand N, Campanella M, Campos-Salinas J, Candi E, Cao L, Caplan AB, Carding SR, Cardoso SM, Carew JS, Carlin CR, Carmignac V, Carneiro LA, Carra S, Caruso RA, Casari G, Casas C, Castino R, Cebollero E, Cecconi F, Celli J, Chaachouay H, Chae HJ, Chai CY, Chan DC, Chan EY, Chang RC, Che CM, Chen CC, Chen GC, Chen GQ, Chen M, Chen Q, Chen SS, Chen W, Chen X, Chen YG, Chen Y, Chen YJ, Chen Z, Cheng A, Cheng CH, Cheng Y, Cheong H, Cheong JH, Cherry S, Chess-Williams R, Cheung ZH, Chevet E, Chiang HL, Chiarelli R, Chiba T, Chin LS, Chiou SH, Chisari FV, Cho CH, Cho DH, Choi AM, Choi D, Choi KS, Choi ME, Chouaib S, Choubey D, Choubey V, Chu CT, Chuang TH, Chueh SH, Chun T, Chwae YJ, Chye ML, Ciarcia R, Ciriolo MR, Clague MJ, Clark RS, Clarke PG, Clarke R, Codogno P, Coller HA, Colombo MI, Comincini S, Condello M, Condorelli F, Cookson MR, Coombs GH, Coppens I, Corbalan R, Cossart P, Costelli P, Costes S, Coto-Montes A, Couve E, Coxon FP, Cregg JM, Crespo JL, Cronje MJ, Cuervo AM, Cullen JJ, Czaja MJ, D'Amelio M, Darfeuille-Michaud A, Davids LM, Davies FE, De Felici M, de Groot JF, de Haan CA, De Martino L, De Milito A, De Tata V, Debnath J, Degterev A, Dehay B, Delbridge LM, Demarchi F, Deng YZ, Dengjel J, Dent P, Denton D, Deretic V, Desai SD, Devenish RJ, Di Gioacchino M, Di Paolo G, Di Pietro C, Diaz-Araya G, Diaz-Laviada I, Diaz-Meco MT, Diaz-Nido J, Dikic I, Dinesh-Kumar SP, Ding WX, Distelhorst CW, Diwan A, Djavaheri-Mergny M, Dokudovskaya S, Dong Z, Dorsey FC, Dosenko V, Dowling JJ, Doxsey S, Dreux M, Drew ME, Duan Q, Duchosal MA, Duff K, Dugail I, Durbeej M, Duszenko M, Edelstein CL, Edinger AL, Egea G, Eichinger L, Eissa NT, Ekmekcioglu S, El-Deiry WS, Elazar Z, Elgendy M, Ellerby LM, Eng KE, Engelbrecht AM, Engelender S, Erenpreisa J, Escalante R, Esclatine A, Eskelinen EL, Espert L, Espina V, Fan H, Fan J, Fan QW, Fan Z, Fang S, Fang Y, Fanto M, Fanzani A, Farkas T, Farre JC, Faure M, Fechheimer M, Feng CG, Feng J, Feng Q, Feng Y, Fesus L, Feuer R, Figueiredo-Pereira ME, Fimia GM, Fingar DC, Finkbeiner S, Finkel T, Finley KD, Fiorito F, Fisher EA, Fisher PB, Flajolet M, Florez-McClure ML, Florio S, Fon EA, Fornai F, Fortunato F, Fotedar R, Fowler DH, Fox HS, Franco R, Frankel LB, Fransen M, Fuentes JM, Fueyo J, Fujii J, Fujisaki K, Fujita E, Fukuda M, Furukawa RH, Gaestel M, Gailly P, Gajewska M, Galliot B, Galy V, Ganesh S, Ganetzky B, Ganley IG, Gao FB, Gao GF, Gao J, Garcia L, Garcia-Manero G, Garcia-Marcos M, Garmyn M, Gartel AL, Gatti E, Gautel M, Gawriluk TR, Gegg ME, Geng J, Germain M, Gestwicki JE, Gewirtz DA, Ghavami S, Ghosh P, Giammarioli AM, Giromanolaki AN, Gibson SB, Gilkerson RW, Ginger ML, Ginsberg HN, Golab J, Goligorsky MS, Golstein P, Gomez-Manzano C, Goncu E, Gongora C, Gonzalez CD, Gonzalez R, Gonzalez-Estevez C, Gonzalez-Polo RA, Gonzalez-Rey E, Gorbunov NV, Gorski S, Goruppi S, Gottlieb RA, Gozuacik D, Granato GE, Grant GD, Green KN, Gregorc A, Gros F, Grose C, Grunt TW, Gual P, Guan JL, Guan KL, Guichard SM, Gukovskaya AS, Gukovskiy I, Gunst J, Gustafsson AB, Halayko AJ, Hale AN, Halonen SK, Hamasaki M, Han F, Han T, Hancock MK, Hansen M, Harada H, Harada M, Hardt SE, Harper JW, Harris AL, Harris J, Harris SD, Hashimoto M, Haspel JA, Hayashi S, Hazelhurst LA, He C, He YW, Hebert MJ, Heidenreich KA, Helfrich MH, Helgason GV, Henske EP, Herman B, Herman PK, Hetz C, Hilfiker S, Hill JA, Hocking LJ, Hofman P, Hofmann TG, Hohfeld J, Holyoake TL, Hong MH, Hood DA, Hotamisligil GS, Houwerzijl EJ, Hoyer-Hansen M, Hu B, Hu CA, Hu HM, Hua Y, Huang C, Huang J, Huang S, Huang WP, Huber TB, Huh WK, Hung TH, Hupp TR, Hur GM, Hurley JB, Hussain SN, Hussey PJ, Hwang JJ, Hwang S, Ichihara A, Ilkhanizadeh S, Inoki K, Into T, Iovane V, Iovanna JL, Ip NY, Isaka Y, Ishida H, Isidoro C, Isobe K, Iwasaki A, Izquierdo M, Izumi Y, Jaakkola PM, Jaattela M, Jackson GR, Jackson WT, Janji B, Jendrach M, Jeon JH, Jeung EB, Jiang H, Jiang JX, Jiang M, Jiang Q, Jiang X, Jimenez A, Jin M, Jin S, Joe CO, Johansen T, Johnson DE, Johnson GV, Jones NL, Joseph B, Joseph SK, Joubert AM, Juhasz G, Juillerat-Jeanneret L, Jung CH, Jung YK, Kaarniranta

K, Kaasik A, Kabuta T, Kadowaki M, Kagedal K, Kamada Y, Kaminsky VO, Kampinga HH, Kanamori H, Kang C, Kang KB, Kang KI, Kang R, Kang YA, Kanki T, Kanneganti TD, Kanno H, Kanthasamy AG, Kanthasamy A, Karantza V, Kaushal GP, Kaushik S, Kawazoe Y, Ke PY, Kehrl JH, Kelekar A, Kerkhoff C, Kessel DH, Khalil H, Kiel JA, Kiger AA, Kihara A, Kim DR, Kim DH, Kim EK, Kim HR, Kim JS, Kim JH, Kim JC, Kim JK, Kim PK, Kim SW, Kim YS, Kim Y, Kimchi A, Kimmelman AC, King JS, Kinsella TJ, Kirkin V, Kirshenbaum LA, Kitamoto K, Kitazato K, Klein L, Klimecki WT, Klucken J, Knecht E, Ko BC, Koch JC, Koga H, Koh JY, Koh YH, Koike M, Komatsu M, Kominami E, Kong HJ, Kong WJ, Korolchuk VI, Kotake Y, Koukourakis MI, Kouri Flores JB, Kovacs AL, Kraft C, Krainc D, Kramer H, Kretz-Remy C, Krichevsky AM, Kroemer G, Kruger R, Krut O, Ktistakis NT, Kuan CY, Kucharczyk R, Kumar A, Kumar R, Kumar S, Kundu M, Kung HJ, Kurz T, Kwon HJ, La Spada AR, Lafont F, Lamark T, Landry J, Lane JD, Lapayette P, Laporte JF, Laszlo L, Lavandero S, Lavoie JN, Layfield R, Lazo PA, Le W, Le Cam L, Ledbetter DJ, Lee AJ, Lee BW, Lee GM, Lee J, Lee JH, Lee M, Lee MS, Lee SH, Leeuwenburgh C, Legembre P, Legouis R, Lehmann M, Lei HY, Lei QY, Leib DA, Leiro J, Lemasters JJ, Lemoine A, Lesniak MS, Lev D, Levenson VV, Levine B, Levy E, Li F, Li JL, Li L, Li S, Li W, Li XJ, Li YB, Li YP, Liang C, Liang Q, Liao YF, Liberski PP, Lieberman A, Lim HJ, Lim KL, Lim K, Lin CF, Lin FC, Lin J, Lin JD, Lin K, Lin WW, Lin WC, Lin YL, Linden R, Lingor P, Lippincott-Schwartz J, Lisanti MP, Liton PB, Liu B, Liu CF, Liu K, Liu L, Liu QA, Liu W, Liu YC, Liu Y, Lockshin RA, Lok CN, Lonial S, Loos B, Lopez-Berestein G, Lopez-Otin C, Lossi L, Lotze MT, Low P, Lu B, Lu Z, Luciano F, Lukacs NW, Lund AH, Lynch-Day MA, Ma Y, Macian F, MacKeigan JP, Macleod KF, Madeo F, Maiuri L, Maiuri MC, Malagoli D, Malicdan MC, Malorni W, Man N, Mandelkow EM, Manon S, Manov I, Mao K, Mao X, Mao Z, Marambaud P, Marazziti D, Marcel YL, Marchbank K, Marchetti P, Marciniak SJ, Marcondes M, Mardi M, Marfe G, Marino G, Markaki M, Marten MR, Martin SJ, Martinand-Mari C, Martinet W, Martinez-Vicente M, Masini M, Matarrese P, Matsuo S, Matteoni R, Mayer A, Mazure NM, McConkey DJ, McConnell MJ, McDermott C, McDonald C, McInerney GM, McKenna SL, McLaughlin B, McLean PJ, McMaster CR, McQuibban GA, Meijer AJ, Meisler MH, Melendez A, Melia TJ, Melino G, Mena MA, Menendez JA, Menna-Barreto RF, Menon MB, Menzies FM, Mercer CA, Merighi A, Merry DE, Meschini S, Meyer CG, Meyer TF, Miao CY, Miao JY, Michels PA, Michiels C, Mijaljica D, Milojkovic A, Minucci S, Miracco C, Miranti CK, Mitroulis I, Miyazawa K, Mizushima N, Mograbi B, Mohseni S, Molero X, Mollereau B, Mollinedo F, Momoi T, Monastyrska I, Monick MM, Monteiro MJ, Moore MN, Mora R, Moreau K, Moreira PI, Moriyasu Y, Moscat J, Mostowy S, Motttram JC, Motyl T, Moussa CE, Muller S, Munger K, Munz C, Murphy LO, Murphy ME, Musaro A, Mysorekar I, Nagata E, Nagata K, Nahimana A, Nair U, Nakagawa T, Nakahira K, Nakano H, Nakatogawa H, Nanjundan M, Naqvi NI, Narendra DP, Narita M, Navarro M, Nawrocki ST, Nazarko TY, Nemchenko A, Netea MG, Neufeld TP, Ney PA, Nezis IP, Nguyen HP, Nie D, Nishino I, Nislow C, Nixon RA, Noda T, Noegel AA, Nogalska A, Noguchi S, Notterpek L, Novak I, Nozaki T, Nukina N, Nurnberger T, Nyfeler B, Obara K, Oberley TD, Oddo S, Ogawa M, Ohashi T, Okamoto K, Oleinick NL, Oliver FJ, Olsen LJ, Olsson S, Opota O, Osborne TF, Ostrand GK, Otsu K, Ou JH, Ouimet M, Overholzer M, Ozpolat B, Paganetti P, Pagnini U, Pallet N, Palmer GE, Palumbo C, Pan T, Panaretakis T, Pandey UB, Papackova Z, Papassideri I, Paris I, Park J, Park OK, Parys JB, Parzych KR, Patschan S, Patterson C, Pattingre S, Pawelek JM, Peng J, Perlmutter DH, Perrotta I, Perry G, Pervaiz S, Peter M, Peters GJ, Petersen M, Petrovski G, Phang JM, Piacentini M, Pierre P, Pierrefite-Carle V, Pierron G, Pinkas-Kramarski R, Piras A, Piri N, Platanias LC, Poggeler S, Poirot M, Poletti A, Pous C, Pozuelo-Rubio M, Praetorius-Ibba M, Prasad A, Prescott M, Priault M, Produit-Zengaffinen N, Progulske-Fox A, Proikas-Cezanne T, Przedborski S, Przyklenk K, Puertollano R, Puyal J, Qian SB, Qin L, Qin ZH, Quaggin SE, Raben N, Rabinowich H, Rabkin SW, Rahman I, Rami A, Ramm G, Randall K, Randow F, Rao VA, Rathmell JC, Ravikumar B, Ray SK, Reed BH, Reed JH, Reggiori F, Regnier-Vigouroux A, Reichert AS, Reiners JJ, Jr., Reiter RJ, Ren J, Revuelta JL, Rhodes CJ, Ritis K, Rizzo E, Robbins J, Roberge M, Roca H, Roccheri MC, Rocchi S, Rodemann HP, Rodriguez de Cordoba S, Rohrer B, Roninson IB, Rosen K, Rost-Roszkowska MM, Rouis M, Rouschop KM, Rovetta F, Rubin BP, Rubinsztein DC, Ruckdeschel K, Rucker EB, 3rd, Rudich A, Rudolf E, Ruiz-Opazo N, Russo R, Rusten TE, Ryan KM, Ryter SW, Sabatini DM, Sadoshima J, Saha T, Saitoh T, Sakagami H, Sakai Y, Salekdeh GH, Salomoni P, Salvaterra PM, Salvesen G, Salvioli R, Sanchez AM, Sanchez-Alcazar JA, Sanchez-Prieto R, Sandri M, Sankar U, Sansanwal P, Santambrogio L, Saran S, Sarkar S, Sarwal M, Sasakawa C, Sasnauskiene A, Sass M, Sato K, Sato M, Schapira AH, Scharl M, Schatzl HM, Scheper W, Schiaffino S, Schneider C, Schneider ME, Schneider-Stock R, Schoenlein PV, Schorderet DF, Schuller C, Schwartz GK, Scorrano L, Sealy L, Seglen PO, Segura-Aguilar J, Seiliez I, Seleverstov O, Sell C, Seo JB, Separovic D, Setaluri V, Setoguchi T, Settembre C, Shacka JJ, Shanmugam M, Shapiro IM, Shaulian E, Shaw RJ, Shelhamer JH, Shen HM, Shen WC, Sheng ZH, Shi Y, Shibuya K, Shidoji Y, Shieh JJ, Shih CM, Shimada Y, Shimizu S, Shintani T, Shirihai OS, Shore GC, Sibirny AA, Sidhu SB, Sikorska B, Silva-Zacarin EC, Simmons A, Simon AK, Simon HU, Simone C, Simonsen A, Sinclair DA, Singh R, Sinha D, Sinicrope FA, Sirko A, Siu PM, Sivridis E, Skop V, Skulachev VP, Slack RS, Smaili SS, Smith DR, Soengas MS, Soldati T, Song X, Sood AK, Soong TW, Sotgia F, Spector SA, Spies CD, Springer W, Srinivasula SM, Stefanis L, Steffan JS, Stendel R, Stenmark H, Stephanou A, Stern ST, Sternberg C, Stork B, Stralfors P, Subauste CS, Sui X, Sulzer D, Sun J, Sun SY, Sun ZJ, Sung JJ, Suzuki K, Suzuki T, Swanson MS, Swanton C, Sweeney ST, Sy LK, Szabadkai G, Tabas I, Taegtmeier H, Tafani M, Takacs-Vellai K, Takano Y, Takegawa K, Takemura G, Takeshita F, Talbot NJ, Tan KS, Tanaka K, Tang D, Tanida I, Tannous BA, Tavernarakis N, Taylor GS, Taylor GA, Taylor JP, Terada LS, Terman A, Tettamanti G, Thevisen K, Thompson CB, Thorburn A, Thumm M, Tian F, Tian Y, Tocchini-Valentini G, Tolkovsky AM, Tomino Y, Tonges L, Tooze SA, Tournier C, Tower J, Towns R, Trajkovic V, Travassos LH, Tsai TF, Tschann MP, Tsubata T, Tsung A, Turk B, Turner LS, Tyagi SC,

- Uchiyama Y, Ueno T, Umekawa M, Umemiya-Shirafuji R, Unni VK, Vaccaro MI, Valente EM, Van den Berghe G, van der Klei IJ, van Doorn W, van Dyk LF, van Egmond M, van Grunsven LA, Vandenabeele P, Vandenberghe WP, Vanhorebeek I, Vaquero EC, Velasco G, Vellai T, Vicencio JM, Vierstra RD, Vila M, Vindis C, Viola G, Viscomi MT, Voitsekhovskaja OV, von Haefen C, Votruba M, Wada K, Wade-Martins R, Walker CL, Walsh CM, Walter J, Wan XB, Wang A, Wang C, Wang D, Wang F, Wang G, Wang H, Wang HG, Wang HD, Wang J, Wang K, Wang M, Wang RC, Wang X, Wang YJ, Wang Y, Wang Z, Wang ZC, Wansink DG, Ward DM, Watada H, Waters SL, Webster P, Wei L, Weihl CC, Weiss WA, Welford SM, Wen LP, Whitehouse CA, Whitton JL, Whitworth AJ, Wileman T, Wiley JW, Wilkinson S, Willbold D, Williams RL, Williamson PR, Wouters BG, Wu C, Wu DC, Wu WK, Wytenbach A, Xavier RJ, Xi Z, Xia P, Xiao G, Xie Z, Xu DZ, Xu J, Xu L, Xu X, Yamamoto A, Yamashina S, Yamashita M, Yan X, Yanagida M, Yang DS, Yang E, Yang JM, Yang SY, Yang W, Yang WY, Yang Z, Yao MC, Yao TP, Yeganeh B, Yen WL, Yin JJ, Yin XM, Yoo OJ, Yoon G, Yoon SY, Yorimitsu T, Yoshikawa Y, Yoshimori T, Yoshimoto K, You HJ, Youle RJ, Younes A, Yu L, Yu SW, Yu WH, Yuan ZM, Yue Z, Yun CH, Yuzaki M, Zabirnyk O, Silva-Zacarin E, Zacks D, Zacksenhaus E, Zaffaroni N, Zakeri Z, Zeh HJ, 3rd, Zeitlin SO, Zhang H, Zhang HL, Zhang J, Zhang JP, Zhang L, Zhang MY, Zhang XD, Zhao M, Zhao YF, Zhao Y, Zhao ZJ, Zheng X, Zhivotovsky B, Zhong Q, Zhou CZ, Zhu C, Zhu WG, Zhu XF, Zhu X, Zhu Y, Zoladek T, Zong WX, Zorzano A, Zschocke J, and Zuckerbraun B. Guidelines for the use and interpretation of assays for monitoring autophagy. *Autophagy* 8: 445–544, 2012.
96. Kornmann B. The molecular hug between the ER and the mitochondria. *Curr Opin Cell Biol* 25: 443–448, 2013.
97. Kornmann B, Currie E, Collins SR, Schuldiner M, Nunnari J, Weissman JS, and Walter P. An ER-mitochondria tethering complex revealed by a synthetic biology screen. *Science* 325: 477–481, 2009.
98. Kroemer G, Galluzzi L, and Brenner C. Mitochondrial membrane permeabilization in cell death. *Physiol Rev* 87: 99–163, 2007.
99. Kuwana T, Mackey MR, Perkins G, Ellisman MH, Latterich M, Schneider R, Green DR, and Newmeyer DD. Bid, Bax, and lipids cooperate to form supramolecular openings in the outer mitochondrial membrane. *Cell* 111: 331–342, 2002.
100. Le S, Connors TJ, and Maroney AC. c-Jun N-terminal kinase specifically phosphorylates p66ShcA at serine 36 in response to ultraviolet irradiation. *J Biol Chem* 276: 48332–48336, 2001.
101. Lebedzinska M, Duszynski J, Rizzuto R, Pinton P, and Wieckowski MR. Age-related changes in levels of p66Shc and serine 36-phosphorylated p66Shc in organs and mouse tissues. *Arch Biochem Biophys* 486: 73–80, 2009.
102. Lebedzinska M, Karkucinska-Wieckowska A, Giorgi C, Karczmarewicz E, Pronicka E, Pinton P, Duszynski J, Pronicki M, and Wieckowski MR. Oxidative stress-dependent p66Shc phosphorylation in skin fibroblasts of children with mitochondrial disorders. *Biochim Biophys Acta* 1797: 952–960, 2010.
103. Lebedzinska M, Karkucinska-Wieckowska A, Wojtala A, Suski JM, Szabadkai G, Wilczynski G, Wlodarczyk J, Diogo CV, Oliveira PJ, Tauber J, Jezek P, Pronicki M, Duszynski J, Pinton P, and Wieckowski MR. Disrupted ATP synthase activity and mitochondrial hyperpolarisation-dependent oxidative stress is associated with p66Shc phosphorylation in fibroblasts of NARP patients. *Int J Biochem Cell Biol* 45: 141–150, 2013.
104. Lebedzinska M, Szabadkai G, Jones AW, Duszynski J, and Wieckowski MR. Interactions between the endoplasmic reticulum, mitochondria, plasma membrane and other subcellular organelles. *Int J Biochem Cell Biol* 41: 1805–1816, 2009.
105. Lee GS, Subramanian N, Kim AI, Aksentijevich I, Goldbach-Mansky R, Sacks DB, Germain RN, Kastner DL, and Chae JJ. The calcium-sensing receptor regulates the NLRP3 inflammasome through Ca^{2+} and cAMP. *Nature* 492: 123–127, 2012.
106. Lee SY, Hwang DY, Kim YK, Lee JW, Shin IC, Oh KW, Lee MK, Lim JS, Yoon DY, Hwang SJ, and Hong JT. PS2 mutation increases neuronal cell vulnerability to neurotoxicants through activation of caspase-3 by enhancing of ryanodine receptor-mediated calcium release. *FASEB J* 20: 151–153, 2006.
107. Lee YJ, Jeong SY, Karbowski M, Smith CL, and Youle RJ. Roles of the mammalian mitochondrial fission and fusion mediators Fis1, Drp1, and Opa1 in apoptosis. *Mol Biol Cell* 15: 5001–5011, 2004.
108. Leissring MA, Parker I, and LaFerla FM. Presenilin-2 mutations modulate amplitude and kinetics of inositol 1, 4,5-trisphosphate-mediated calcium signals. *J Biol Chem* 274: 32535–32538, 1999.
109. Li G, Mongillo M, Chin KT, Harding H, Ron D, Marks AR, and Tabas I. Role of ERO1- α -mediated stimulation of inositol 1,4,5-trisphosphate receptor activity in endoplasmic reticulum stress-induced apoptosis. *J Cell Biol* 186: 783–792, 2009.
110. Li M, Chiou KR, and Kass DA. Shear stress inhibition of H₂O₂ induced p66(Shc) phosphorylation by ASK1-JNK inactivation in endothelium. *Heart Vessels* 22: 423–427, 2007.
111. Li N, Ragheb K, Lawler G, Sturgis J, Rajwa B, Melendez JA, and Robinson JP. Mitochondrial complex I inhibitor rotenone induces apoptosis through enhancing mitochondrial reactive oxygen species production. *J Biol Chem* 278: 8516–8525, 2003.
112. Ly JD, Grubb DR, and Lawen A. The mitochondrial membrane potential ($\Delta\psi(m)$) in apoptosis; an update. *Apoptosis* 8: 115–128, 2003.
113. Marchi S, Lupini L, Patergnani S, Rimessi A, Missiroli S, Bonora M, Bononi A, Corra F, Giorgi C, De Marchi E, Poletti F, Gafa R, Lanza G, Negrini M, Rizzuto R, and Pinton P. Downregulation of the mitochondrial calcium uniporter by cancer-related miR-25. *Curr Biol* 23: 58–63, 2013.
114. Marchi S, Patergnani S, and Pinton P. The endoplasmic reticulum-mitochondria connection: one touch, multiple functions. *Biochim Biophys Acta* 1837: 461–469, 2014.
115. Marchi S and Pinton P. The mitochondrial calcium uniporter complex: molecular components, structure and pathophysiological implications. *J Physiol* 592: 829–839, 2014.
116. Marchi S, Rimessi A, Giorgi C, Baldini C, Ferroni L, Rizzuto R, and Pinton P. Akt kinase reducing endoplasmic reticulum Ca^{2+} release protects cells from Ca^{2+} -dependent apoptotic stimuli. *Biochem Biophys Res Commun* 375: 501–505, 2008.
117. Mariathasan S, Weiss DS, Newton K, McBride J, O'Rourke K, Roose-Girma M, Lee WP, Weinrauch Y, Monack DM, and Dixit VM. Cryopyrin activates the in-

- flammasome in response to toxins and ATP. *Nature* 440: 228–232, 2006.
118. Marongiu R, Spencer B, Crews L, Adame A, Patrick C, Trejo M, Dallapiccola B, Valente EM, and Masliah E. Mutant Pink1 induces mitochondrial dysfunction in a neuronal cell model of Parkinson's disease by disturbing calcium flux. *J Neurochem* 108: 1561–1574, 2009.
 119. Martin LJ, Pan Y, Price AC, Sterling W, Copeland NG, Jenkins NA, Price DL, and Lee MK. Parkinson's disease alpha-synuclein transgenic mice develop neuronal mitochondrial degeneration and cell death. *J Neurosci* 26: 41–50, 2006.
 120. May D, Itin A, Gal O, Kalinski H, Feinstein E, and Keshet E. Ero1-L alpha plays a key role in a HIF-1-mediated pathway to improve disulfide bond formation and VEGF secretion under hypoxia: implication for cancer. *Oncogene* 24: 1011–1020, 2005.
 121. Migliaccio E, Giorgio M, and Pelicci PG. Apoptosis and aging: role of p66Shc redox protein. *Antioxid Redox Signal* 8: 600–608, 2006.
 122. Migliaccio E, Mele S, Salcini AE, Pelicci G, Lai KM, Superti-Furga G, Pawson T, Di Fiore PP, Lanfrancione L, and Pelicci PG. Opposite effects of the p52shc/p46shc and p66shc splicing isoforms on the EGF receptor-MAP kinase-fos signalling pathway. *EMBO J* 16: 706–716, 1997.
 123. Murakami T, Ockinger J, Yu J, Byles V, McColl A, Hofer AM, and Horng T. Critical role for calcium mobilization in activation of the NLRP3 inflammasome. *Proc Natl Acad Sci U S A* 109: 11282–11287, 2012.
 124. Nakahira K, Haspel JA, Rathinam VA, Lee SJ, Dolinay T, Lam HC, Englert JA, Rabinovitch M, Cernadas M, Kim HP, Fitzgerald KA, Ryter SW, and Choi AM. Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome. *Nat Immunol* 12: 222–230, 2011.
 125. Naon D and Scorrano L. At the right distance: ER-mitochondria juxtaposition in cell life and death. *Biochim Biophys Acta* 1843: 2184–2194, 2014.
 126. Napoli C, Martin-Padura I, de Nigris F, Giorgio M, Mansueti G, Somma P, Condorelli M, Sica G, De Rosa G, and Pelicci P. Deletion of the p66Shc longevity gene reduces systemic and tissue oxidative stress, vascular cell apoptosis, and early atherogenesis in mice fed a high-fat diet. *Proc Natl Acad Sci U S A* 100: 2112–2116, 2003.
 127. Narendra D, Tanaka A, Suen DF, and Youle RJ. Parkin is recruited selectively to impaired mitochondria and promotes their autophagy. *J Cell Biol* 183: 795–803, 2008.
 128. Nauseef WM. Detection of superoxide anion and hydrogen peroxide production by cellular NADPH oxidases. *Biochim Biophys Acta* 1840: 757–767, 2014.
 129. Nemoto S, Combs CA, French S, Ahn BH, Fergusson MM, Balaban RS, and Finkel T. The mammalian longevity-associated gene product p66shc regulates mitochondrial metabolism. *J Biol Chem* 281: 10555–10560, 2006.
 130. Oelze M, Kroller-Schon S, Steven S, Lubos E, Doppler C, Hausding M, Tobias S, Brochhausen C, Li H, Torzewski M, Wenzel P, Bachschmid M, Lackner KJ, Schulz E, Munzel T, and Daiber A. Glutathione peroxidase-1 deficiency potentiates dysregulatory modifications of endothelial nitric oxide synthase and vascular dysfunction in aging. *Hypertension* 63: 390–396, 2014.
 131. Okada S, Kao AW, Ceresa BP, Blaikie P, Margolis B, and Pessin JE. The 66-kDa Shc isoform is a negative regulator of the epidermal growth factor-stimulated mitogen-activated protein kinase pathway. *J Biol Chem* 272: 28042–28049, 1997.
 132. Olichon A, Baricault L, Gas N, Guillou E, Valette A, Belenguer P, and Lenaers G. Loss of OPA1 perturbs the mitochondrial inner membrane structure and integrity, leading to cytochrome c release and apoptosis. *J Biol Chem* 278: 7743–7746, 2003.
 133. Orsini F, Migliaccio E, Moroni M, Contursi C, Raker VA, Piccini D, Martin-Padura I, Pelliccia G, Trinei M, Bono M, Puri C, Tacchetti C, Ferrini M, Mannucci R, Nicoletti I, Lanfrancione L, Giorgio M, and Pelicci PG. The life span determinant p66Shc localizes to mitochondria where it associates with mitochondrial heat shock protein 70 and regulates trans-membrane potential. *J Biol Chem* 279: 25689–25695, 2004.
 134. Osowski CM, Hara T, O'Sullivan-Murphy B, Kanekura K, Lu S, Hara M, Ishigaki S, Zhu LJ, Hayashi E, Hui ST, Greiner D, Kaufman RJ, Bortell R, and Urano F. Thiorodoxin-interacting protein mediates ER stress-induced beta cell death through initiation of the inflammasome. *Cell Metab* 16: 265–273, 2012.
 135. Ottolini D, Cali T, Negro A, and Brini M. The Parkinson disease-related protein DJ-1 counteracts mitochondrial impairment induced by the tumour suppressor protein p53 by enhancing endoplasmic reticulum-mitochondria tethering. *Hum Mol Genet* 22: 2152–2168, 2013.
 136. Panov AV, Gutekunst CA, Leavitt BR, Hayden MR, Burke JR, Strittmatter WJ, and Greenamyre JT. Early mitochondrial calcium defects in Huntington's disease are a direct effect of polyglutamines. *Nat Neurosci* 5: 731–736, 2002.
 137. Papanicolaou KN, Khairallah RJ, Ngoh GA, Chikando A, Luptak I, O'Shea KM, Riley DD, Lugus JJ, Colucci WS, Lederer WJ, Stanley WC, and Walsh K. Mitofusin-2 maintains mitochondrial structure and contributes to stress-induced permeability transition in cardiac myocytes. *Mol Cell Biol* 31: 1309–1328, 2011.
 138. Paradies G, Petrosillo G, Paradies V, and Ruggiero FM. Role of cardiolipin peroxidation and Ca²⁺ in mitochondrial dysfunction and disease. *Cell Calcium* 45: 643–650, 2009.
 139. Park J, Lee SB, Lee S, Kim Y, Song S, Kim S, Bae E, Kim J, Shong M, Kim JM, and Chung J. Mitochondrial dysfunction in Drosophila PINK1 mutants is complemented by parkin. *Nature* 441: 1157–1161, 2006.
 140. Park S, Juliana C, Hong S, Datta P, Hwang I, Fernandes-Alnemri T, Yu JW, and Alnemri ES. The mitochondrial antiviral protein MAVS associates with NLRP3 and regulates its inflammasome activity. *J Immunol* 191: 4358–4366, 2013.
 141. Patergnani S, Marchi S, Rimessi A, Bonora M, Giorgi C, Mehta KD, and Pinton P. PRKCB/protein kinase C, beta and the mitochondrial axis as key regulators of autophagy. *Autophagy* 9: 1367–1385, 2013.
 142. Patergnani S, Suski JM, Agnoletto C, Bononi A, Bonora M, De Marchi E, Giorgi C, Marchi S, Missiroli S, Poletti F, Rimessi A, Duszynski J, Wieckowski MR, and Pinton P. Calcium signaling around Mitochondria Associated Membranes (MAMs). *Cell Commun Signal* 9: 19, 2011.
 143. Petrosillo G, Colantuono G, Moro N, Ruggiero FM, Tiravanti E, Di Venosa N, Fiore T, and Paradies G. Melatonin protects against heart ischemia-reperfusion injury by inhibiting mitochondrial permeability transition pore opening. *Am J Physiol Heart Circ Physiol* 297: H1487–H1493, 2009.

144. Pinton P, Giorgi C, and Pandolfi PP. The role of PML in the control of apoptotic cell fate: a new key player at ER-mitochondria sites. *Cell Death Differ* 18: 1450–1456, 2011.
145. Pinton P, Giorgi C, Siviero R, Zecchini E, and Rizzuto R. Calcium and apoptosis: ER-mitochondria Ca^{2+} transfer in the control of apoptosis. *Oncogene* 27: 6407–6418, 2008.
146. Pinton P, Rimessi A, Marchi S, Orsini F, Migliaccio E, Giorgio M, Contursi C, Minucci S, Mantovani F, Wieckowski MR, Del Sal G, Pelicci PG, and Rizzuto R. Protein kinase C beta and prolyl isomerase 1 regulate mitochondrial effects of the life-span determinant p66Shc. *Science* 315: 659–663, 2007.
147. Poston CN, Krishnan SC, and Bazemore-Walker CR. In-depth proteomic analysis of mammalian mitochondria-associated membranes (MAM). *J Proteomics* 79: 219–230, 2013.
148. Prause J, Goswami A, Katona I, Roos A, Schnizler M, Bushuven E, Dreier A, Buchkremer S, Johann S, Beyer C, Deschauer M, Troost D, and Weis J. Altered localization, abnormal modification and loss of function of Sigma receptor-1 in amyotrophic lateral sclerosis. *Hum Mol Genet* 22: 1581–1600, 2013.
149. Raturi A and Simmen T. Where the endoplasmic reticulum and the mitochondrion tie the knot: the mitochondria-associated membrane (MAM). *Biochim Biophys Acta* 1833: 213–224, 2013.
150. Reboredo M, Greaves RF, and Hahn G. Human cytomegalovirus proteins encoded by UL37 exon 1 protect infected fibroblasts against virus-induced apoptosis and are required for efficient virus replication. *J Gen Virol* 85: 3555–3567, 2004.
151. Redmann M, Dodson M, Boyer-Guittaut M, Darley-Usmar V, and Zhang J. Mitophagy mechanisms and role in human diseases. *Int J Biochem Cell Biol* 53: 127–133, 2014.
152. Reichert AS and Neupert W. Contact sites between the outer and inner membrane of mitochondria-role in protein transport. *Biochim Biophys Acta* 1592: 41–49, 2002.
153. Rimessi A, Marchi S, Patergnani S, and Pinton P. H-Ras-driven tumoral maintenance is sustained through caveolin-1-dependent alterations in calcium signaling. *Oncogene* 33: 2329–2340, 2014.
154. Rizzuto R, Pinton P, Carrington W, Fay FS, Fogarty KE, Lifshitz LM, Tuft RA, and Pozzan T. Close contacts with the endoplasmic reticulum as determinants of mitochondrial Ca^{2+} responses. *Science* 280: 1763–1766, 1998.
155. Rossol M, Pierer M, Raulien N, Quandt D, Meusch U, Rothe K, Schubert K, Schoneberg T, Schaefer M, Krugel U, Smajilovic S, Brauner-Osborne H, Baerwald C, and Wagner U. Extracellular Ca^{2+} is a danger signal activating the NLRP3 inflammasome through G protein-coupled calcium sensing receptors. *Nat Commun* 3: 1329, 2012.
156. Sano R, Annunziata I, Patterson A, Moshiah S, Gomero E, Opferman J, Forte M, and d'Azzo A. GM1-ganglioside accumulation at the mitochondria-associated ER membranes links ER stress to Ca^{2+} -dependent mitochondrial apoptosis. *Mol Cell* 36: 500–511, 2009.
157. Saxena G, Chen J, and Shalev A. Intracellular shuttling and mitochondrial function of thioredoxin-interacting protein. *J Biol Chem* 285: 3997–4005, 2010.
158. Schon EA and Area-Gomez E. Mitochondria-associated ER membranes in Alzheimer disease. *Mol Cell Neurosci* 55: 26–36, 2013.
159. Schroder K and Tschopp J. The inflammasomes. *Cell* 140: 821–832, 2010.
160. Seth RB, Sun L, Ea CK, and Chen ZJ. Identification and characterization of MAVS, a mitochondrial antiviral signaling protein that activates NF-kappaB and IRF 3. *Cell* 122: 669–682, 2005.
161. Shavali S, Brown-Borg HM, Ebadi M, and Porter J. Mitochondrial localization of alpha-synuclein protein in alpha-synuclein overexpressing cells. *Neurosci Lett* 439: 125–128, 2008.
162. Shimizu S, Matsuoka Y, Shinohara Y, Yoneda Y, and Tsujimoto Y. Essential role of voltage-dependent anion channel in various forms of apoptosis in mammalian cells. *J Cell Biol* 152: 237–250, 2001.
163. Sies H. Role of metabolic H₂O₂ generation: redox signaling and oxidative stress. *J Biol Chem* 289: 8735–8741, 2014.
164. Simmen T, Aslan JE, Blagoveshchenskaya AD, Thomas L, Wan L, Xiang Y, Feliciangeli SF, Hung CH, Crump CM, and Thomas G. PACS-2 controls endoplasmic reticulum-mitochondria communication and Bid-mediated apoptosis. *EMBO J* 24: 717–729, 2005.
165. Simmen T, Lynes EM, Gesson K, and Thomas G. Oxidative protein folding in the endoplasmic reticulum: tight links to the mitochondria-associated membrane (MAM). *Biochim Biophys Acta* 1798: 1465–1473, 2010.
166. Spescha RD, Glanzmann M, Simic B, Witassek F, Keller S, Akhmedov A, Tanner FC, Luscher TF, and Camici GG. Adaptor protein p66(Shc) mediates hypertension-associated, cyclic stretch-dependent, endothelial damage. *Hypertension* 64: 347–353, 2014.
167. Subramanian N, Natarajan K, Clatworthy MR, Wang Z, and Germain RN. The adaptor MAVS promotes NLRP3 mitochondrial localization and inflammasome activation. *Cell* 153: 348–361, 2013.
168. Suski JM, Lebieczinska M, Wojtala A, Duszynski J, Giorgi C, Pinton P, and Wieckowski MR. Isolation of plasma membrane-associated membranes from rat liver. *Nat Protoc* 9: 312–322, 2014.
169. Szalai G, Krishnamurthy R, and Hajnoczky G. Apoptosis driven by IP(3)-linked mitochondrial calcium signals. *EMBO J* 18: 6349–6361, 1999.
170. Trinei M, Giorgio M, Cicalese A, Barozzi S, Ventura A, Migliaccio E, Milia E, Padura IM, Raker VA, Maccarana M, Petronilli V, Minucci S, Bernardi P, Lanfrancone L, and Pelicci PG. A p53-p66Shc signalling pathway controls intracellular redox status, levels of oxidation-damaged DNA and oxidative stress-induced apoptosis. *Oncogene* 21: 3872–3878, 2002.
171. Tschopp J. Mitochondria: Sovereign of inflammation? *Eur J Immunol* 41: 1196–1202, 2011.
172. Tu BP and Weissman JS. Oxidative protein folding in eukaryotes: mechanisms and consequences. *J Cell Biol* 164: 341–346, 2004.
173. Tu H, Nelson O, Bezprozvanny A, Wang Z, Lee SF, Hao YH, Serneels L, De Strooper B, Yu G, and Bezprozvanny I. Presenilins form ER Ca^{2+} leak channels, a function disrupted by familial Alzheimer's disease-linked mutations. *Cell* 126: 981–993, 2006.
174. Verfaillie T, Rubio N, Garg AD, Bultynck G, Rizzuto R, Decuyper JP, Piette J, Linehan C, Gupta S, Samali A, and Agostinis P. PERK is required at the ER-mitochondrial contact sites to convey apoptosis after ROS-based ER stress. *Cell Death Differ* 19: 1880–1891, 2012.
175. Wang X, Winter D, Ashrafi G, Schlehe J, Wong YL, Selkoe D, Rice S, Steen J, LaVoie MJ, and Schwarz TL. PINK1 and Parkin target Miro for phosphorylation and

- degradation to arrest mitochondrial motility. *Cell* 147: 893–906, 2011.
176. Wardman P. Fluorescent and luminescent probes for measurement of oxidative and nitrosative species in cells and tissues: progress, pitfalls, and prospects. *Free Radic Biol Med* 43: 995–1022, 2007.
177. Wasiak S, Zunino R, and McBride HM. Bax/Bak promote sumoylation of DRP1 and its stable association with mitochondria during apoptotic cell death. *J Cell Biol* 177: 439–450, 2007.
178. Wenzel P, Knorr M, Kossmann S, Stratmann J, Hausding M, Schuhmacher S, Karbach SH, Schwenk M, Yogev N, Schulz E, Oelze M, Grabbe S, Jonuleit H, Becker C, Daiber A, Waisman A, and Munzel T. Lysozyme M-positive monocytes mediate angiotensin II-induced arterial hypertension and vascular dysfunction. *Circulation* 124: 1370–1381, 2011.
179. Wieckowski MR, Giorgi C, Lebedzinska M, Duszynski J, and Pinton P. Isolation of mitochondria-associated membranes and mitochondria from animal tissues and cells. *Nat Protoc* 4: 1582–1590, 2009.
180. Wiley SE, Andreyev AY, Divakaruni AS, Karisch R, Perkins G, Wall EA, van der Geer P, Chen YF, Tsai TF, Simon MI, Neel BG, Dixon JE, and Murphy AN. Wolfram Syndrome protein, Miner1, regulates sulphhydryl redox status, the unfolded protein response, and Ca²⁺ homeostasis. *EMBO Mol Med* 5: 904–918, 2013.
181. Winterbourn CC. The challenges of using fluorescent probes to detect and quantify specific reactive oxygen species in living cells. *Biochim Biophys Acta* 1840: 730–738, 2014.
182. Winther JR and Thorpe C. Quantification of thiols and disulfides. *Biochim Biophys Acta* 1840: 838–846, 2014.
183. Wood-Kaczmar A, Gandhi S, Yao Z, Abramov AY, Miljan EA, Keen G, Stanyer L, Hargreaves I, Klupsch K, Deas E, Downward J, Mansfield L, Jat P, Taylor J, Heales S, Duchon MR, Latchman D, Tabrizi SJ, and Wood NW. PINK1 is necessary for long term survival and mitochondrial function in human dopaminergic neurons. *PLoS One* 3: e2455, 2008.
184. Yang Y, Gehrke S, Imai Y, Huang Z, Ouyang Y, Wang JW, Yang L, Beal MF, Vogel H, and Lu B. Mitochondrial pathology and muscle and dopaminergic neuron degeneration caused by inactivation of Drosophila Pink1 is rescued by Parkin. *Proc Natl Acad Sci U S A* 103: 10793–10798, 2006.
185. Yu W, Sun Y, Guo S, and Lu B. The PINK1/Parkin pathway regulates mitochondrial dynamics and function in mammalian hippocampal and dopaminergic neurons. *Hum Mol Genet* 20: 3227–3240, 2011.
186. Zaid H, Abu-Hamad S, Israelson A, Nathan I, and Shoshan-Barmatz V. The voltage-dependent anion channel-1 modulates apoptotic cell death. *Cell Death Differ* 12: 751–760, 2005.
187. Zatti G, Ghidoni R, Barbiero L, Binetti G, Pozzan T, Fasolato C, and Pizzo P. The presenilin 2 M239I mutation associated with familial Alzheimer's disease reduces Ca²⁺ release from intracellular stores. *Neurobiol Dis* 15: 269–278, 2004.
188. Zhang A, Williamson CD, Wong DS, Bullough MD, Brown KJ, Hathout Y, and Colberg-Poley AM. Quantitative proteomic analyses of human cytomegalovirus-induced restructuring of endoplasmic reticulum-mitochondrial contacts at late times of infection. *Mol Cell Proteomics* 10: M111 009936, 2011.
189. Zhou R, Tardivel A, Thorens B, Choi I, and Tschopp J. Thioredoxin-interacting protein links oxidative stress to inflammasome activation. *Nat Immunol* 11: 136–140, 2010.
190. Zhou R, Yazdi AS, Menu P, and Tschopp J. A role for mitochondria in NLRP3 inflammasome activation. *Nature* 469: 221–225, 2011.

Address correspondence to:

Dr. Paolo Pinton
Section of Pathology, Oncology
and Experimental Biology
Laboratory for Technologies
of Advanced Therapies (LTTA)
Department of Morphology, Surgery
and Experimental Medicine
University of Ferrara
Ferrara 44100
Italy

E-mail: pnp@unife.it

Date of first submission to ARS Central, December 10, 2014;
date of acceptance, January 1, 2015.

Abbreviations Used

Ψ_m = the mitochondrial membrane potential
 $A\beta$ = amyloid- β
 AD = Alzheimer's disease
 AIF = apoptosis-inducing factor
 ALS = amyotrophic lateral sclerosis
 ASC = apoptosis-associated speck-like protein
 containing a caspase recruitment domain
 ASK1 = apoptosis signal-regulating kinase 1
 CaSR = murine calcium-sensing receptor
 CL = cardiolipin
 CNX = calnexin
 DRP1 = dynamin-related protein 1
 ER = endoplasmic reticulum
 ERAD = ER-associated degradation
 ERK = extracellular signal-regulated kinase
 ERp44 = endoplasmic reticulum resident protein 44
 FAD = familial Alzheimer's disease
 FATP4 = fatty acid transport protein 4
 GM1 = GM1-ganglioside
 GRP75 = glucose-regulated protein 75-kDa
 HIV-1 = human immunodeficiency virus type 1
 IMM = inner mitochondrial membrane
 IMS = mitochondrial intermembrane space
 IP3R = inositol 1,4,5-trisphosphate receptor
 JNK = c-Jun N-terminal kinase
 LRRs = leucine-rich repeats
 MAMs = mitochondria-associated membranes
 MAVS = mitochondrial antiviral signaling protein
 MCU = mitochondrial calcium uniporter
 MDA5 = melanoma differentiation-associated gene 5
 MMP = mitochondrial outer membrane permeabilization
 mPTP = mitochondrial permeability transition pore
 OMM = outer mitochondrial membrane
 PACS-2 = phosphofurin acidic cluster sorting protein 2
 PAMs = plasma membranes associated membranes

Abbreviations Used (Cont.)

PD = Parkinson's disease
PDI = protein disulfide-isomerase
PEMT2 = phosphatidylethanolamine
N-methyltransferase 2
PKC β = protein kinase C β
PML = promyelocytic leukemia protein
PP2A = phosphatase A2
PS1 = presenilin-1
PS2 = presenilin-2

ROS = reactive oxygen species
SERAC1 = serine active site containing 1
SIG1R = sigma-1 receptor
STING = stimulator of interferon genes
TXNIP = thioredoxin interacting protein
UPR = unfolded protein response
VAPB = vesicle-associated membrane protein-associated
protein B
VDAC = voltage-dependent anion channel
Vpr = viral protein R

This article has been cited by:

1. Yin Fei, Cadenas Enrique. 2015. Mitochondria: The Cellular Hub of the Dynamic Coordinated Network. *Antioxidants & Redox Signaling* **22**:12, 961-964. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]