

Novel role for polycystin-1 in modulating cell proliferation through calcium oscillations in kidney cells

G. Aguiari*, V. Trimi*, M. Bogo*, A. Mangolini*, G. Szabadkai†, P. Pinton†, R. Witzgall§, P. C. Harris¶, P. A. Borea‡, R. Rizzuto† and L. del Senno*

*Department of Biochemistry and Molecular Biology; †General Pathology, Department of Experimental and Diagnostic Medicine; ‡Pharmacology Unit, Department of Clinical and Experimental Medicine, University of Ferrara, Ferrara, Italy, §Institute for Molecular and Cellular Anatomy, University of Regensburg, Regensburg, Germany, and ¶Division of Nephrology and Hypertension, Mayo Clinic College of Medicine, Rochester, MN, USA

Received 17 July 2007; revision accepted 18 October 2007

OnlineOpen: This article is available free online at www.blackwell-synergy.com

Abstract. *Objectives:* Polycystin-1 (PC1), a signalling receptor regulating Ca^{2+} -permeable cation channels, is mutated in autosomal dominant polycystic kidney disease, which is typically characterized by increased cell proliferation. However, the precise mechanisms by which PC1 functions on Ca^{2+} homeostasis, signalling and cell proliferation remain unclear. Here, we investigated the possible role of PC1 as a modulator of non-capacitative Ca^{2+} entry (NCCE) and Ca^{2+} oscillations, with downstream effects on cell proliferation. *Results and discussion:* By employing RNA interference, we show that depletion of endogenous PC1 in HEK293 cells leads to an increase in serum-induced Ca^{2+} oscillations, triggering nuclear factor of activated T cell activation and leading to cell cycle progression. Consistently, Ca^{2+} oscillations and cell proliferation are increased in PC1-mutated kidney cystic cell lines, but both abnormal features are reduced in cells that exogenously express PC1. Notably, blockers of the NCCE pathway, but not of the CCE, blunt abnormal oscillation and cell proliferation. Our study therefore provides the first demonstration that PC1 modulates Ca^{2+} oscillations and a molecular mechanism to explain the association between abnormal Ca^{2+} homeostasis and cell proliferation in autosomal dominant polycystic kidney disease.

INTRODUCTION

Increases in cytosolic concentration of Ca^{2+} ions $[\text{Ca}^{2+}]_i$ regulate virtually all cellular processes, including events such as survival, proliferation and apoptosis, which decide the fate of each cell (Berridge *et al.* 2000). These functions are disturbed in autosomal dominant polycystic kidney disease (ADPKD), one of the most common inherited diseases in humans (1 in 500 to 1 in 1000), and one that accounts for about 10% of end-stage renal failure. The primary phenotype

Correspondence: Laura del Senno, Department of Biochemistry and Molecular Biology, University of Ferrara, Ferrara, Italy. Tel.: 39 532 455447; Fax: 39 532 202723; E-mail: sen@unife.it

Re-use of this article is permitted in accordance with the Creative Commons Deed, Attribution 2.5, which does not permit commercial exploitation.

of ADPKD is the progressive expansion, in both kidneys, of multiple fluid-filled cysts, which gradually replace normal renal tissue and lead to end-stage renal failure and dialysis dependency by mid-to-late adulthood (Gabow 1990). The systemic nature of the disease is determined by a number of extrarenal manifestations (from hepatic and brain cysts to heart valve and cerebrovascular abnormalities) that accompany the kidney cysts (Gabow 1990) and indicate the complex role of ADPKD causative gene products (Ong & Harris 2005).

In approximately 85% of ADPKD cases, the disease arises as a consequence of loss of function and missense mutations of the *PKD1* gene, which encodes polycystin-1 (PC1), with the remaining cases being caused by mutations in polycystin-2-encoding *PKD2* gene (PC2) (Mochizuki *et al.* 1996; Harris 1999). The phenotypes associated with either genotype are otherwise very similar, indicating that the two proteins function with the same pathways. PC1 and PC2 are the prototype members of a new family of integral membrane proteins that play important roles in a variety of biological processes, including fertilization, ion translocation and mechanotransduction (Delmas 2004). PC1 (around 450 kDa and 11 transmembrane domains) acts as an orphan receptor mainly involved in control of epithelial cell population growth (Bhunia *et al.* 2002; Kim *et al.* 2004; Li *et al.* 2005; Manzati *et al.* 2005), migration (Nickel *et al.* 2002; Polgar *et al.* 2005), differentiation (Aguari *et al.* 1998; Puri *et al.* 2004) and apoptosis (Boca *et al.* 2006). In association with PC2, a Ca^{2+} -regulated Ca^{2+} -permeable non-selective cation channel (Hanaoka *et al.* 2000; González-Perrett *et al.* 2001; Koulou *et al.* 2002), PC1 is required for regulation of the cell cycle (Bhunia *et al.* 2002) and activation of cation permeable currents (Vandorpe *et al.* 2001; Babich *et al.* 2004; Pelucchi *et al.* 2006); it is also thought to be involved in regulation of G-protein signalling (Parnell *et al.* 1998; Delmas *et al.* 2002).

Our group has recently found that expression of a membrane-targeted PC1 C-terminus in kidney HEK293 cells increases cell response to ATP-evoked $[Ca^{2+}]_i$ (Aguari *et al.* 2003) and serum-evoked cell proliferation (Manzati *et al.* 2005). It has also recently been reported that the treatment of PKD-cystic cells with molecules causing sustained increases in $[Ca^{2+}]_i$ may reverse their mitogenic response (Yamaguchi *et al.* 2006). Therefore, the aim of the present work was to determine the mechanism by which PC1 modulates Ca^{2+} levels, considering the possibility that a reduction in endogenous PC1 by small-interference RNA might affect Ca^{2+} mobilization and thus induce a growth-stimulated phenotype characteristic of PKD cells. Our results, obtained by employing pharmacological tools on kidney cell lines depleted of endogenous PC1 by PKD1 RNA interference or in PKD1 mutated cystic cell lines, demonstrate that down-regulation of PC1 leads to elevation in Ca^{2+} oscillations by stimulating non-capacitative Ca^{2+} entry (NCCE). Such oscillatory signals trigger nuclear factor of activated T cell (NFAT) activation and other Ca^{2+} -dependent signalling pathways, leading to increased cell proliferation. Furthermore, the inhibitory effects of exogenously expressed PC1 on Ca^{2+} oscillations, as well as proliferation of HEK293 cells, indicate PC1 as a physiological modulator of NCCE, which plays a crucial role on basal and evoked $[Ca^{2+}]_i$ levels. These findings pave the way to future investigations on involvement of such channels in both the physiology and pathophysiology of kidney cells.

MATERIALS AND METHODS

Reagents

Dulbecco's modified Eagle's medium/F12 and minimum essential medium media, G418 antibiotic, bovine serum albumin (BSA), 1-(beta-[3-(4-methoxyphenyl) propoxy]-4-methoxyphenethyl)-1H-imidazole hydrochloride (SKF96365), cyclosporin A, gadolinium and anti-FLAG M2 mouse

monoclonal antibody were obtained from Sigma-Aldrich (Milano, Italy), foetal bovine serum (FBS) was obtained from Eurobio (Celbio, Milan, Italy) and selective cell-permeable inhibitors of protein kinase C (PKC)- α and PKC- β 1 (Ro-320432 and hispidin, respectively), were purchased from Calbiochem (La Jolla, CA, USA) and protease inhibitors were from Roche Diagnostics (Monza, Italy). Rabbit polyclonal antitubulin and antirabbit rhodamine- and FITC-conjugated antibody were obtained from Santa Cruz (DBA Italia Srl, Segrate, Italy), while rabbit polyclonal antip27 antibody was obtained from Cell Signalling Technology (Celbio SRL, Italy). Enhanced chemiluminescent substrates for Western blotting (SuperSignalDura or SuperSignalFemto) and horseradish peroxidase-conjugated goat antirabbit and antimouse antibodies were purchased from Pierce (Celbio S.r.l., Milan, Italy). The pSUPER RNAi System was obtained from OligoEngine (Seattle, WA, USA). (R,S)-(3,4-dihydro-6,7-dimethoxy-isochinolin-1-yl)-2-phenyl-N,N-di[2-(2,3,4-trimethoxyphenyl)ethyl] acetamid mesylate (LOE908) was kindly provided by Boehringer Ingelheim (Ingelheim, Germany).

Production of siRNA expressing vectors

Two siRNAs (*a* and *b*) for PC1 were constructed according to a published method (Brummelkamp *et al.* 2002). Briefly, 64 nt primers were designed to include a 19-nt PKD1 sequence, its complement, a spacer region, 5' *Bg*III site and 3' *Hind*III site.

The forward PKD1a primer sequence: 5'-GATCCCCCGACAAGCAGTCCCTGACCTTC-AAGAGAGGTCAGGGACTGCTTGTCGTTTTTAAA-3' and the reverse primer sequence: 5'-AGCTTTTCCAAAAACGACAAGCAGTCCCTGACCTCTCTTGAAGGTCAGGGACTG-CTTGTCCGGG-3'.

The forward PKD1b primer sequence: 5'-GATCCCCCTTACGCGGAGCGGCACGT-TCAAGAGACGTGCCGCTCCGCGTGAAGTTTTTGGAAA-3' and the reverse primer sequence: 5'-AGCTTTTCCAAAACTTACGCGGAGCGGCACGTCTCTTGAACGTGCC-GCTCCGCGTGAAGGGG-3'. The target PKD1 sequence was substituted by a scramble nucleotide sequence as control: 5'-TTCTCCGAACGTGTCACGT-3'. Annealed double strands of DNAs were cloned into the pSUPER plasmid vector following the Oligoengine procedure.

Cells, stable transfection, cell cycle and proliferation assays

HEK293 cells were maintained and cultured as previously described (Manzati *et al.* 2005). Wild-type (pSUPER) and recombinant constructs (pSiScramble, pSsiPKD1 *a* and *b*) were co-transfected with pCDNA3 (ratio 1 : 10, respectively) by the calcium phosphate method, in HEK293 cells (Pelucchi *et al.* 2006), and G418-resistant clones (HEK293^{pSUPER}, HEK293^{pSiScramble} and HEK293^{pSsiPKD1}), and were screened by Western blotting analysis, through detection of endogenous PC1 protein using 7e12 monoclonal antibody (Qian *et al.* 2003). PKD1 RNA levels were analysed by real-time RT-PCR, as previously reported (Aguiari *et al.* 2004). PKD1 RNA quantification was obtained by the Δ CT method. Transcription of the ACTIN housekeeping gene was used as endogenous RNA control for normalization and the Δ CT was then calculated (Δ CT = CT(target gene) – CT(actin gene)). Final results, expressed as *N*fold content in PKD1 gene expression relative to the ACTIN gene, were determined as follows: $N(\text{target}) = 2^{-\Delta\Delta\text{CT}}$, as previously described (Aguiari *et al.* 2004).

Cell cycle analysis was done with the use of flow cytometry of propidium iodide-treated cells, using the FACSCalibur Becton Dickinson Immunocytometry System. Previously described SV40-transformed human cell lines (one derived from normal and two from cystic kidney cell lines) (Nauli *et al.* 2006), were grown in Dulbeccos's modified Eagle's medium/F12, 10% FBS. Cystic cells carry the Q2556X PKD1 mutation: one is heterozygous, while the other is hemizygous, for deletion of the normal allele (Nauli *et al.* 2006). After plating at low density

(30 000 cells/mL) in 24-well plates, cell proliferation was measured by direct cell counting (Manzati *et al.* 2005), and by 24 and 48 h incubation with [3H] thymidine (1 μ Ci/mL).

Transient transfection and immunofluorescence analysis

HEK293^{pSUPER} and HEK293^{pSipKD1} cells were cultured on 24-mm coverslips for 24 h then were transiently transfected with full-length FLAG-tagged mouse PC1 (Grimm *et al.* 2003) by Ca^{2+} phosphate precipitation. After 3 days of transfection, non-fixed cells were washed twice with phosphate-buffered saline (PBS) buffer and incubated at room temperature for 1 h with anti-FLAG M2 mouse monoclonal antibody at a dilution of 1 : 200, in a solution containing 0.2% gelatin. After three washes with PBS, cells were treated with a 0.2% gelatin PBS solution that contained secondary antirabbit rhodamine-conjugated antibody at room temperature in the dark for 1 h. After three washes in PBS, cells were analysed using a Zeiss Axiovert 200 fluorescence microscope equipped with a back-illuminated CCD camera (Roper Scientific, Tucson, AZ, USA), excitation and emission filter wheels (Sutter Instrument Company, Novato, CA, USA), and piezoelectric motoring of the Z stage (Physik Instrumente, GmbH and Co., Karlsruhe, Germany).

Ca^{2+} measurement

Measurements of FBS-evoked calcium levels were performed in cells grown on coverslips and transfected with recombinant cytoplasmic aequorin cDNA (Aguiari *et al.* 2003). Seventy-two hours after aequorin reconstitution, cells on coverslips were transferred into the perfusion chamber of a luminometer and, following 1 min of perfusion, stimulated with 1% FBS. The relationship between photon emission (L/L_{max}) and Ca^{2+} concentration allows conversion by computer algorithm of light released into the specimen. L and L_{max} are the instant and maximal rates of light emission, the former having been obtained after 1% FBS stimulation and the latter brought about by cell lysis with 100 μ M digitonin (Manzati *et al.* 2005).

Cytoplasmic Ca^{2+} oscillations were measured in Fura-2AM-loaded cells (Manzati *et al.* 2005). Cells, grown on coverslips, were starved ON with BSA, loaded with Fura-2AM (4 μ M/30 min/37 °C), and transferred to the thermostated stage of a Zeiss Axiovert 200 inverted microscope equipped with a Sutter filterwheel and 340/380 excitation filters. Cells were stimulated either with 1% FBS alone or with other molecules for 15 min or 30 min, acquiring 1 image every 1 s or 3 s, respectively. Signals were computed into relative ratio units of the fluorescence intensity of different wavelengths (340/380 nm). Ca^{2+} oscillation amplitude was calculated by measuring Ca^{2+} concentration in Fura-2AM-loaded cells, as previously described (Manzati *et al.* 2005). Calibration was carried out using MetaFluor software, according to the equation $[Ca^{2+}] = K_d \times (Sf2/Sb2) \times (R - R_{min}) / (R_{max} - R)$, using $K_d = 224$ nM. R_{max} and R_{min} were determined at the end of each experiment in KRB/ Ca^{2+} /1 μ M ionomycin, and KRB/1 mM EGTA/ionomycin, respectively.

NFAT and PKC- α fluorescence analysis

Cells were transfected with either NFAT-green fluorescent protein (GFP) or PKC- α -GFP constructs by using the Ca^{2+} phosphate procedure (Manzati *et al.* 2005). NFAT and PKC- α subcellular localization was analysed using a digital fluorescence microscope, and images were captured and recorded using a digital-imaging system (Manzati *et al.* 2005).

NFAT luciferase assay

After 24 h plating in a 6-well plastic plate, cells were transiently transfected with 4 μ g of pNFAT-TA-Luc plasmid DNA by the Ca^{2+} phosphate method, and 6 h post-transfection, they were washed and incubated for an additional 20 h in medium supplemented with 1% FBS. After cell treatment with passive lysis buffer (Promega, Madison, WI, USA), 25 μ l of cell lysate was added

to 50 μ l of luciferase substrate (Promega) and then the samples were analysed using a 20/20⁰ luminometer (Turner Biosystems, Sunnyvale, CA, USA).

Statistical analysis

Where applicable, statistical errors of averaged data were given as means \pm SD or SEM and statistical significance was assessed by Student's *t*-test.

RESULTS

Relationship between PC1 depletion by RNA interference and proliferation of HEK293 cells

A variable, but consistent down-regulation of PC1 was accomplished in HEK293^{pSsiPKD1} cell clones obtained by stable transfection with pSUPER plasmid expressing PKD1 siRNAs (Fig. 1a). An average of more than ~50% reduction in PC1 protein was found in clones transfected by either the *a* or *b* PKD1 siRNA-cloned sequence, as compared to those transfected with the empty vector or that expressing the scramble nucleotide sequence, as controls [Fig. 1b, PC1/ β -tubulin: 0.70 ± 0.09 and 0.55 ± 0.5 in HEK293^{pSsiPKD1} *a* and *b* clones (pSsiPKD1), respectively, versus 1.52 ± 0.11 and 1.46 ± 0.34 in control HEK293^{pSUPER} and HEK293^{pSiScramble} clones (pS and pS^{Sc}); $P < 0.01$ and $P < 0.05$, respectively, versus pS and $P < 0.05$ versus pSc]. This inhibition of PC1 by siRNA was found to be selective as neither variations in β -tubulin were observed, nor PC1 levels altered, in clones expressing the control vectors. Moreover, extent of PC1 inhibition was consistent with the reduction in PKD1 RNA, as shown in HEK293^{pSsiPKD1} *a* cells by real-time RT-PCR analysis (Fig. 1c).

Because the loss of PC1 in kidney cysts is associated with proliferation of epithelial cells, and HEK293 cells overexpressing the putative dominant negative PC1 C-terminus have been shown to increase cell proliferation (Manzati *et al.* 2005), HEK293^{pSsiPKD1} stably transfected cells were assessed for possible changes in cell proliferation. PC1 inhibition did indeed cause a significant increase in 1% FBS-induced cell proliferation, as illustrated by the examples shown in Fig. 1d. This increase was detectable both in [³H] thymidine uptake after 24 and 48 h incubation, and after 3 and 5 days of culture ($106\,667 \pm 5700$ versus $62\,333 \pm 8500$ and $65\,100 \pm 11\,300$ after 3 days of culture, and $203\,333 \pm 6500$ versus $126\,667 \pm 5600$ and $120\,600 \pm 17\,800$ after 5 days, respectively, in HEK293^{pSsiPKD1} cells versus HEK293^{pSUPER} and HEK293^{pSiScramble} control cells; $P < 0.01$ and $P < 0.001$ after 3 and 5 days, respectively). Furthermore, proliferation increase in HEK^{pSsiPKD1} *a*₁ and *a*₃ cells was found to be smaller than that observed in *b*₃ cells, which showed the lowest PC1 levels (Fig. 1e), thus indicating a relationship between degree of PC1 suppression and cell proliferation.

PC1 expression reduces calcium oscillation

The effect of PC1 silencing on $[Ca^{2+}]_i$ evoked by FBS, at 1% concentration used in cell proliferation studies, was initially investigated in HEK^{pSsiPKD1} clone *a*₁ through both aequorin and Fura-2AM experiments.

After transfection with Ca²⁺-sensitive photoprotein aequorin, a slight, although non-significant, increase in serum-induced cytoplasmic Ca²⁺ peak was found in HEK^{pSsiPKD1} cells, as compared to control cells (bars of Fig. 2a). Nevertheless, it is apparent from representative traces of Fig. 2a that Ca²⁺ levels were more sustained and oscillating in HEK^{pSsiPKD1} cells. These differences were confirmed in single-cell Ca²⁺-imaging experiments (Fig. 2b). Representative traces from




Figure 1. PKD1 gene silencing increases cell proliferation in HEK293 cells. (a) Western blot analysis of PC1 from plasma membranes of HEK293 cells stably transfected with plasmids expressing two PKD1-specific siRNA (pSsiPKD1 *a* and *b*) and one scramble RNA sequence (pSsiScramble, pSc), and transfected with the pSUPER empty vector (pS). A representative entire blot of the control HEK293^{pSUPER} clone 1 (pS), and HEK293^{pSsiPKD1} clone a₁ (pSsiPKD1) is included. The analysis of PC1 probed with the monoclonal antibody 7e12 (Qian *et al.* 2003) was related to β -tubulin. The variations in β -tubulin levels were due to unequal protein loading. Two *a* and three *b* HEK293^{pSsiPKD1} clones are shown. (b) The expression of either pSsiPKD1 *a* or *b* construct significantly reduced PC1 production in comparison with that of β -tubulin. Bars show the average relative PC1 expression (\pm SEM) in six controls (three HEK293^{pSUPER}, pS, white bar; and three HEK293^{pSsiScramble}, pSc, grained bar), 10 HEK293^{pSsiPKD1} *a* and 3 HEK293^{pSsiPKD1} *b* clones (dark and light grey bars) from three independent experiments (*a* and *b* versus pS, $**P < 0.01$ and $*P < 0.05$, respectively, unpaired Student's *t*-test). (c) Fold decrease in PKD1 mRNA levels in HEK293^{pSsiPKD1} clone *a* (grey bar) compared to HEK293^{pSUPER} cells (pS, white bar), as detected by real-time RT-PCR analysis. PKD1 RNA levels relative to those of actin RNA were measured by $\Delta\Delta C_t$ values that were calculated as described in the Methods section. Bars show the average (\pm SD) of four different experiments ($**P < 0.01$, unpaired Student's *t*-test). (d) [³H] thymidine incorporation after both 24 and 48 h incubation with 1% FBS was higher in HEK293^{pSsiPKD1} clone b₃ (grey bar) than to HEK293^{pSUPER} control cells (white bar); moreover, HEK293^{pSsiPKD1} cell proliferation was also increased after 3 and 5 days of culture in the presence of 1% FBS (HEK293^{pSUPER} cells, white bars, HEK293^{pSsiScramble} cells, grained bars, and HEK293^{pSsiPKD1} clone a₂, grey bars, the only one studied after 5 days). The data shown are the mean (\pm SD) of at least two independent experiments ($*P < 0.05$, $**P < 0.01$ and $***P < 0.001$, unpaired Student's *t*-test). (e) After 3 days, cell proliferation increase was greater in HEK293^{pSsiPKD1} b₃ clone showing lower PC1 expression (hatched bars).

HEK293^{pSUPER} and HEK293^{pSsiPKD1} cells revealed a pattern of changes in FBS-evoked Ca^{2+} levels that was characterized by oscillation frequency that was consistently higher in HEK293^{pSsiPKD1} cells. The oscillation increase was observed in 81.5% of cells ($n = 76$). When the analysis was performed in different HEK^{pSsiPKD1} clones, mean oscillation frequency increased ~ 2 -fold as compared to control cells (9.78 versus 5.52 and 5.36 in HEK293^{pSsiPKD1} and HEK^{pSsiScramble} cells, respectively, the mean values in a 15-min period, $P < 0.01$) (Fig. 2c), suggesting a relationship between reduced PC1 expression and increased Ca^{2+} oscillation. Oscillation amplitude, expressed as Ca^{2+} concentration, was not different in HEK293^{pSUPER} and HEK293^{pSsiPKD1} cells (98.05 ± 29.12 nM and 102.00 ± 20.89 nM in HEK293^{pSUPER} and HEK293^{pSsiPKD1} cells, respectively, Fig. 2d), lending further support to the hypothesized association between PC1 deficiency and oscillation frequency. Because the difference in oscillation frequency at 10% FBS concentration was markedly reduced (from about 2 in 1% FBS to about 1.4-fold in 10% FBS, Fig. 2e), further experiments were performed in 1% FBS. Moreover, because no significant differences were




Figure 2. PKD1 gene silencing increases FBS-induced Ca^{2+} oscillation frequency in HEK293 cells. (a) Representative cytoplasmic Ca^{2+} levels from control HEK293^{pSUPER} 1 (pS, blue trace) and HEK293^{pSsiPKD1} a₁ (red trace) cells. Ca^{2+} levels were measured in cells transfected with the Ca^{2+} -binding photoprotein cytAEQ. Where indicated, cells were challenged with 1% FBS. Bars represent the peak evoked by FBS stimulation (averages ± SEM, $P = 0.057$) in control (blue) and HEK293^{pSsiPKD1} (red) clones. (b) FBS increased Ca^{2+} oscillation frequency in PKD1-specific siRNA expressing cells. Representative Ca^{2+} signals from an individual control pS (blue trace) and HEK293^{pSsiPKD1} a₁ cell (red trace) loaded with Fura-2-AM and stimulated with 1% FBS. Average distribution of Ca^{2+} oscillation frequency (± SD, white parts of the bars) from HEK293^{pSUPER} 1 ($n = 66$ cells, four experiments) and HEK293^{pSsiPKD1} a₁ cells (61 = cells min period, four experiments) in 15 min after FBS stimulation. (c) Bars indicate the average Ca^{2+} oscillation frequency in four HEK293^{pSUPER}, three HEK293^{pSsiScramble} (black bar) and four HEK293^{pSsiPKD1} different clones. The data are the means ± SEM of five experiments ($P < 0.01$ versus pS, unpaired Student's t -test). (d) Average (± SD) of Ca^{2+} oscillation amplitude in three HEK293^{pSUPER} (blue bar) and five HEK293^{pSsiPKD1} (red bar) clones. Amplitude is expressed as Ca^{2+} concentration, measured as described in the Materials and Methods section. (e) Calcium oscillations (average ± SD) evoked by treatment with 0%, 1% and 10% FBS in three HEK293^{pSUPER} (blue bars) and four HEK293^{pSsiPKD1} (red bars) clones. (f) Cell proliferation increase as a function of Ca^{2+} oscillation frequency in four different HEK293^{pSsiPKD1} clones relative to the HEK293^{pSUPER} 1 control clone.

observed between HEK293^{pSUPER} and HEK293^{pSsiScramble} cells, further experiments were mainly performed using HEK293^{pSUPER} cells.

Consistent with increase in proliferation observed in HEK293^{pSsiPKD1} cells, oscillation frequency was higher in HEK293^{pSsiPKD1} clones with higher proliferation rates (Fig. 2f), strongly indicating the relation between Ca^{2+} oscillation frequency and cell proliferation. The b₃ clone, which exhibited greatest oscillation and proliferation associated with the most pronounced PC1 down-regulation (Fig. 1e), was used in the further experiments.

To test the hypothesis of a role for PC1 in regulating Ca^{2+} oscillatory patterns, the latter were examined in HEK293^{pSsiPKD1} cells transiently transfected with a cDNA plasmid expressing mouse full-length PC1-containing N-ter FLAG and C-ter HA tags (Grimm *et al.* 2003). Exogenous expression of mouse PC1 (mPC1) was demonstrated by anti-FLAG immunoblotting in total




Figure 3. Exogenous expression of PC1 reduces Ca²⁺ oscillation frequency in HEK293 cells. (a) Representative Ca²⁺ oscillation patterns from individual HEK293^{pSsiPKD1} (red) and HEK293^{pSUPER} (blue) cells before (upper traces) and after (lower traces) transient transfection with the mouse PC1 expressing plasmid (mPC1). Cells were grown on coverslips, transfected with the plasmid DNA as described in the Materials and Methods section, loaded with Fura-2-AM after 48 h and stimulated with 1% FBS, as described in Fig. 2. Inset: the expression of the mouse PC1 was confirmed by Western blotting and immunofluorescence analysis, as shown in HEK293^{pSUPER} (pS) and HEK293^{pSsiPKD1} (pSsiPKD1) transfected cells, the latter showing that mPC1 is not silenced by human PKD1 siRNA. Cells were lysed and total extracts were analysed by immunoblotting with the anti-FLAG M2 mouse monoclonal antibody recognizing the FLAG-tagged mouse PC1, as described in the Materials and Methods section. The antibody identified a band of about 400 kDa in only mPC1 cDNA-transfected HEK293^{pSUPER} and HEK293^{pSsiPKD1} cells. The ~200 kDa band was deemed to be aspecific as present in all samples. For immunofluorescence analysis, cells were fixed and treated with M2 antibody, as described in the Materials and Methods section. Upper and lower panels: contrast phase and fluorescence images. Staining of plasma membranes was indicated by arrows, mainly at cell–cell interactions. (b) Average percent reduction in Ca²⁺ oscillation frequency in a 15-min period obtained after transient transfection with the mouse PC1 expressing plasmid of HEK293^{pSsiPKD1} clone b₃ (red bars; *n* = 122 cells, five experiments) and HEK293^{pSUPER} cells (blue bars, *n* = 60, three experiments). (c) Reduction in cell proliferation in HEK293 cells stably expressing the full length mouse PC1. pCDNA3 stably transfected control cells (white bar) and cells stably transfected with the full length mouse PKD1 cDNA plasmid (grey bar) were grown for 3 days in presence of 1% FBS. Data are expressed as the average values (± SD; **P* < 0.05 unpaired Student's *t*-test) obtained from two control and four PC1-transfected clones that were plated as described in the Materials and Methods section.

lysates of both HEK293^{pSUPER} and HEK293^{pSsiPKD1} mPC1-transfected cells, and confirmed in at least 20% of cells by anti-FLAG immunofluorescence (inset of Fig. 3), indicating that mPC1 was not silenced by specific human PKD1 siRNA. mPC1 expression caused a clear reduction in oscillation frequency in some cells (Fig. 3a), leading to an average 30% reduction (Fig. 3b, left). Notably, a similar reduction was also observed in PC1-transfected HEK293^{pSUPER} control cells (Fig. 3a,b, right). Combined, these findings suggest that exogenous expression of PC1 is capable of reducing Ca²⁺ oscillation in HEK293 kidney cells. Furthermore, stable transfection of mouse PC1 in HEK293 cells caused a statistically significant reduction in cell proliferation (Fig. 3c). After




Figure 4. Immortalized cyst-lining cells derived from a human kidney with a PKD1 mutation have increased cell proliferation and Ca²⁺ oscillations. (a) Cell cycle analysis in cystic cells, heterozygous (het) and hemizygous (hem) for the Q2556X PKD1 mutation (Nauli *et al.* 2006) showed an increase in S and G₂/M phases compared to normal kidney cells (wt). After 24 h culture in 1% FBS, cells were analysed with propidium iodide staining as described in the Materials and Methods section. (b) Kidney normal (wt) and cystic (het and hem) cells were plated on coverslips 1 day before Ca²⁺ imaging experiments, starved in BSA and loaded with Fura-2-AM as described in the Materials and Methods section. During the experiment, cells were stimulated with 1% FBS and, after the initial Ca²⁺ transient, oscillations were analysed. Representative Ca²⁺ oscillation patterns from individual cells are shown. The average oscillation frequency (\pm SD) in 15 min is indicated by bars (wt, $n = 56$; het, $n = 83$; hem, $n = 51$ cells, in three experiments, $*P < 0.05$ unpaired Student's *t*-test, wt versus hem and versus het).

3 days of 1% FBS treatment, mean cell proliferation was reduced almost by half in four different PC1-transfected clones ($371\ 833 \pm 171\ 660$ versus $766\ 666 \pm 28\ 284$ cells in transfected versus untransfected cells, respectively, $P < 0.05$). This finding demonstrates that PC1-dependent reduction in Ca²⁺ oscillations is strongly associated to reduction in cell proliferation.

Calcium oscillation increases in human PC1-deficient cystic cells

Enhancement of proliferation by PKD1 RNA interference suggests that HEK293^{pSsiPKD1} cells behave in a similar fashion to ADPKD cyst cells, and an increase in cell proliferation and frequency of Ca²⁺ oscillations in cells derived from PKD1 cysts was therefore expected. In order to test this hypothesis immortalized cell lines from kidney cysts, which are heterozygous and putatively hemizygous for the Q2556X PKD1 mutation (Nauli *et al.* 2006), were analysed and results were compared to those obtained with PKD1 wild-type non-cystic cells.

After 1% FBS treatment for 24 h, cell cycle analysis showed that S and G₂/M phases in hemizygous and heterozygous PC1-mutant cystic cells were higher than in wild-type cells, thereby demonstrating a higher proliferation rate of the cystic cells (Fig. 4a). Consistently in these cells, FBS-evoked Ca²⁺ oscillations were increased (Fig. 4b). In particular, average oscillation frequency over a 15-min period was 9.35 ± 1.36 and 7.36 ± 1.16 in cystic versus 3.904 ± 0.04 in normal cells ($P < 0.05$, wild-type versus either hemizygous or heterozygous cells) (bars in Fig. 4b).

Inhibitors of NCCE blunt both increased oscillation and cell proliferation

In order to ascertain the type of channel involved in the increase in FBS-evoked Ca²⁺ oscillations in HEK293^{pSsiPKD1} clones, Ca²⁺ influx was evaluated in the presence of inhibitors of CCE and NCCE (Berridge 2002) (Fig. 5a). The frequency of FBS-evoked Ca²⁺ oscillations in HEK293^{pSsiPKD1} remained unaffected by the presence of 1 μM Gd³⁺, sufficient to block CCE (Mignen *et al.* 2003a). However, oscillations were markedly affected by 5 μM Gd³⁺, which also completely inhibits NCCE (Mignen *et al.* 2003a), in both HEK293^{pSsiPKD1} and HEK293^{pSUPER} control cells. Interestingly, treatment with 100 nM SKF96365, a putative inhibitor of receptor-operated Ca²⁺ entry with NCCE blocker activity at this low concentration (Moneer *et al.* 2003), provoked markedly greater reduction in oscillation frequency in HEK293^{pSsiPKD1} more than in HEK293^{pSUPER} cells. In particular, treatment of HEK293^{pSUPER} cells with either 1 μM Gd³⁺ or 100 nM SKF96365 caused a 38% or 50% reduction in Ca²⁺ oscillations, respectively; conversely,




Figure 5. NCCE-specific inhibitors blunt the effect of the PKD1 gene silencing or PKD1 nonsense mutation on FBS-induced increase in Ca²⁺ oscillations. Effect of gadolinium (Gd³⁺) and SKF96365 on Ca²⁺ oscillation frequency. Representative Ca²⁺ traces from individual HEK293^{pSUPER} (blue lines) and HEK293^{pSsiPKD1} cells (red lines) loaded with Fura-2-AM and treated with the indicated blocker and 1% FBS, are shown. The 1 μM Gd³⁺ did not affect Ca²⁺ oscillations activated by 1% FBS in HEK293^{pSsiPKD1} cells; 5 μM Gd³⁺ and 100 nM SKF96365 markedly reduced Ca²⁺ oscillations. Inset bars represent the average distribution of Ca²⁺ oscillation frequency (± SD) from HEK293^{pSUPER} cells (1 μM Gd³⁺: n = 45 cells; 5 μM Gd³⁺: n = 50 cells; 100 nM SKF96365: n = 52 cells) and from HEK293^{pSsiPKD1} cells (1 μM Gd³⁺: n = 54 cells; 5 μM Gd³⁺: n = 42 cells; 100 nM SKF96365: n = 49 cells) in a 15-min period after FBS stimulation. Results are representative of three independent experiments. Blue and red bars represent percentage reductions (± SD) produced by each treatment in HEK293^{pSUPER} (blue) and HEK293^{pSsiPKD1} cells (red) compared to untreated cells.




Figure 6. NCCE-specific inhibitors blunt the effect of the PKD1 gene silencing or PKD1 nonsense mutation on FBS-induced increase in cell proliferation. (a) The 5 μM Gd³⁺, 100 nM SKF96365 and 30 μM LOE908 inhibitors markedly inhibited the FBS-induced increase in cell proliferation of HEK293^{pSsiPKD1} cells (grey bars) compared to HEK293^{pSUPER} cells (white bars), while 1 μM Gd³⁺ affected to a lesser extent HEK293^{pSsiPKD1} cells. Cells were counted after 48 h culture in 1% FBS, in the absence or presence of the inhibitor. Results obtained in four independent experiments with different clones are shown as means ± SD. In the bar graph (on the right), the blocker effects are expressed in terms of percentage proliferation of untreated cells. (b) Inhibitory effect of NCCE blockers on cell proliferation of cystic kidney cell lines. Cells were treated as reported for HEK293 cells in (a). Only cystic cell proliferation was inhibited by the treatment with the NCCE blockers. Bars represent mean values (± SEM) of three (for SKF96365) and two (for LOE908) independent experiments (**P* < 0.05 unpaired Student's *t*-test) with cells obtained from non-cystic kidney (white), from kidney cysts heterozygous for the Q2556X mutation (light grey) and hemizygous for the same mutation (dark grey).

in HEK293^{pSsiPKD1} cells, 1 μM Gd³⁺ did not affect Ca²⁺ oscillations, while SKF96365 caused an 82% reduction (see percentage values in Fig. 5). Ca²⁺ oscillations were also reduced by 2.5 times in HEK293^{pSsiPKD1} (64%) than in HEK293^{pSUPER} (25%) by treatment with 30 μM LOE908, another NCCE blocker (Moneer & Taylor 2002) (data not shown), suggesting involvement of NCCE activity in abnormal Ca²⁺ oscillations seen in PC1-deficient cells.

Consistently with the results on Ca²⁺ oscillations, 5 μM Gd³⁺, but not 1 μM Gd³⁺, 100 nM SKF96365 and 30 μM LOE908 markedly reduced the FBS-induced cell proliferation in HEK293^{pSsiPKD1} clones (Fig. 6a). As shown in the summary graph of Fig. 6a (showing percentage values), reductions in proliferation caused by different treatments highlighted the strong inhibition caused by NCCE blockers in HEK293^{pSsiPKD1} cell proliferation (67% to 61%), in comparison to the effect in control cells, which was observed to be four times lower (15%).

In order to confirm inhibition also in cyst-derived cells, the effects of NCCE channel blockers were examined in PKD1 Q2556X-mutated cell lines. As expected, cell proliferation was higher in cystic cells than in wild-type cells (152 888.6 ± 18 151 and 130 553.3 ± 10 210 versus 105 777.5 ± 4337, in heterozygous and hemizygous versus wild-type cells, *P* < 0.05) (Fig. 6b). Treatment with either 100 nM SKF96365 or 30 μM LOE908 NCCE blockers did not modify proliferation of wild-type cells, although it did reduce that of PKD1 heterozygous and hemizygous cells (Fig. 6b). Overall, these results confirm the association between abnormal increase in Ca²⁺ oscillations and cell proliferation in ADPKD tubular cells, ascribing the regulatory effect of PC1 mainly to NCCE.

Increase in NFAT activity in HEK293^{pSsiPKD1} cells is associated with cell proliferation

It has been reported that oscillations are an efficient way of differentially coding Ca²⁺ signals and that frequency of oscillations may modulate gene transcription (Dolmetsch *et al.* 1998). NFAT is a Ca²⁺-dependent transcription factor expressed in many cells, including those of the kidney (Puri *et al.* 2004), which is regulated by frequency of Ca²⁺ oscillations (Tomida *et al.* 2003). We, therefore, investigated whether the FBS-induced oscillations observed in HEK293 PKD1-suppressed cells affected NFAT signalling. We monitored NFAT translocation to the nucleus in HEK293^{pSUPER} and HEK293^{pSsiPKD1} cells expressing a GFP-tagged NFAT2. Prior to FBS treatment (0 min of Fig. 7a, left), NFAT-GFP was present in cytoplasmic regions of both HEK293^{pSsiPKD1} and HEK293^{pSUPER} cells, whereas its nuclear translocation following 1% FBS treatment was clearly faster in HEK293^{pSsiPKD1} than in HEK293^{pSUPER} cells. Thus, Ca²⁺ oscillations induced rapid and constant NFAT nuclear translocation in HEK293^{pSsiPKD1}. This faster activation was also observed in PC1-mutant cystic cells (Fig. 7a, right). Moreover, HEK293 cells were




Figure 7. NFAT mediates the cell proliferation increase caused by PKD1 gene silencing and mutation. (a) The fluorescence images of representative cells transfected with NFAT-GFP construct show the distribution of the NFAT-GFP at various time points before and after exposure to 1% FBS. The calcineurin-dependent nuclear translocation of NFAT occurred faster in HEK293^{pSsiPKD1} (pSsiPKD1) and heterozygous (het) cells than in HEK293^{pSUPER} (pS) and in normal kidney (wt) control cells. Arrows show the marked difference in nuclear translocation between silenced and control cells. Pictures are representative of four different experiments. (b) NFAT-dependent transcription was increased by gene silencing of PC1. HEK293^{pSsiPKD1} and control cells were transfected with an NFAT-dependent luciferase reporter plasmid. Cell extracts were collected after 20 h of 1% FBS treatment. Results of luciferase assays from a representative experiment are shown (HEK293^{pSsiPKD1} clone a₁ and b₂, grey bars; HEK293^{pSUPER} control cells, white bar). Results obtained are expressed as fold activation over the luciferase units displayed by control cells. The data are the means ± SEM of five independent experiments (***P* < 0.01 paired Student's *t*-test). (c) Inhibitory effect of 100 nM SKF96365 on luciferase promoter activity in HEK293^{pSsiPKD1} (grey bars) and HEK293^{pSUPER} control cells (white bars). Cells were transfected with an NFAT-dependent luciferase reporter plasmid and results of three independent experiments are expressed as average fold increase in untreated and SKF96365-treated cells. (d) The calcineurin inhibitor Cyclosporin A (CsA) blunted the FBS-induced increase in cell proliferation of HEK293^{pSsiPKD1} cells. Cells were counted after 2 days of culture in 1% FBS. CsA (100 ng/mL) was present or absent as indicated. Results obtained in a triplicate experiment are shown as means ± SD and are representative of four independent experiments obtained from different clones.

transiently transfected with an NFAT-promoter luciferase-reporter plasmid, and luciferase activity was assessed after FBS treatment. Following 2 days of stimulation, ~3.5-fold increase in luciferase activity in HEK293^{pSsiPKD1} cells as compared to HEK293^{pSUPER} cells was observed (Fig. 7b). This suggests that gene expression is modulated by the effect of PC1 depletion on the oscillatory pattern of Ca²⁺ responses and is further supported by the observation that SKF96365 treatment, which blocked HEK293^{pSsiPKD1} cell proliferation, also reduced the NFAT-dependent luciferase activity (Fig. 7c).

In order to test whether the FBS-induced NFAT activation contributed to the increase in proliferation of HEK293^{pSsiPKD1} cells, the cell proliferation rate was analysed in the presence of cyclosporin A (CsA), a typical inhibitor of the Ca²⁺-calmodulin-dependent serine/threonine protein phosphatase calcineurin, which dephosphorylates and activates NFAT (Puri *et al.* 2004). FBS-induced increase in HEK293^{pSsiPKD1} cell proliferation was completely inhibited by CsA (Fig. 7d), supporting the hypothesis of a relationship between the reduction in PC1 levels and increase in both Ca²⁺ oscillations and NFAT-dependent cell proliferation in HEK293^{pSsiPKD1} cells.

Cytoplasmic Ca²⁺ levels and PKC- α activity

Over all, these data suggest a model in which the impairment of PC1 signalling in PC1-deficient cells leads to elevation of serum-evoked Ca²⁺ oscillations, thus activating calcineurin and NFAT signalling and leading ultimately to cell proliferation. This hypothesis is further supported by the measurement of basal Ca²⁺ levels which, in the absence of serum, are higher in HEK293^{pSsiPKD1} than in control cells (140.78 \pm 27.4 versus 85.92 \pm 8.19, respectively). This is consistent with basal Ca²⁺ levels observed in HEK293 cells expressing the isolated cytoplasmic carboxyl terminus of PC1 (Manzati *et al.* 2005). As observed for glutamate-induced intracellular Ca²⁺ oscillations in astrocytes, which involve activation of conventional PKC (Codazzi *et al.* 2001), only serum-induced Ca²⁺ oscillations lead to Ca²⁺ levels sufficient to activate the calcium-dependent PKC- α isoform (Fig. 8a). Serum treatment was indeed accompanied by a prominent recovery of PKC- α -GFP fluorescence in the membrane compartments of HEK293^{pSsiPKD1} cells, while diffuse fluorescence in control cells indicated that the kinase remains confined to cytoplasm.

Because the PKC- α -specific inhibitor Ro-320432, and not the PKC- β inhibitor hispidin, markedly reduced growth of HEK293^{pSsiPKD1} cell population (Fig. 8b); cell proliferation increase was also shown to be dependent on a PKC- α pathway. As we previously observed in HEK293 cells expressing the PC1 C-terminal tail (Manzati *et al.* 2005), in HEK293^{pSsiPKD1} cells this increase was also accompanied by a marked reduction of the growth inhibitory protein p27^{Kip1} (Fig. 8c, upper part). Consistently, p27^{Kip1} was similarly reduced in PKD1 Q2556X-mutated hemizygous cystic cells (Fig. 8c, lower part). Moreover, treatment of HEK293^{pSsiPKD1} cells with either 5 μ M Gd³⁺, 100 nM SKF96365 or 30 μ M LOE908, which reduces cell proliferation, led to increased p27 levels (Fig. 8d).

The present findings, therefore, provide evidence that lack of balance in PC1 levels caused by a reduction in wild-type protein, as well as by PC1 C-tail overexpression, plays a critical role for Ca²⁺ signalling pathways with similar effects on gene expression and subsequently on cell cycle regulation.

DISCUSSION

Our study demonstrates for the first time that PC1 controls Ca²⁺ oscillations and provides a molecular mechanism to explain association of abnormal Ca²⁺ homeostasis and increased cell proliferation in ADPKD, caused by PKD1 mutations. We have shown that either inhibition of




Figure 8. PKD1 gene silencing-dependent cell proliferation is associated to PKC- α activation and reduction of p27^{Kip1} in HEK293 cells. (a) The calcium-dependent membrane translocation of PKC- α occurred in serum treated HEK293^{pSsiPKD1} (pSsiPKD1), but not in control cells (pS). Fluorescence images of cells expressing a PKC- α -GFP construct show that the PKC- α partly translocates to the plasma membrane after 10 min exposure to 1% serum, while control cells maintain a cytoplasm diffuse staining. Arrows indicate the membrane fluorescence. Pictures are representative of three different experiments. (b) The PKC- α -specific inhibition blunted the serum-induced increase in proliferation of HEK293^{pSsiPKD1}. Cells were counted after 48 h culture in 1% FBS in the absence or presence of either a PKC- β -specific (hispidin, 5 μ M) or PKC- α -specific inhibitor (Ro320432, 10 μ M). Results obtained in a triplicate experiment are shown as means \pm SD and are representative of three independent experiments. (c) Whole cell lysates of HEK293^{pSsiPKD1}, HEK293^{pSUPER} cells (upper panel) and of non cystic (wt) and cystic PKD1 Q2556X hemizygous cells (hem) (lower panel), were analysed by immunoblot for expression levels of p27^{Kip1}. The levels were normalized for the expression of actin housekeeping gene. (d) Immunoblot of total lysates of HEK293^{pSUPER} and HEK293^{pSsiPKD1} cells after 24 h treatment with CCE and NCCE blockers. NCCE inhibitors (100 nM SKF96365 and 30 μ M LOE908) markedly increased p27^{Kip1} expression only in HEK293^{pSsiPKD1} cells. Bar graph shows that p27^{Kip1} expression (grey bars) is increased of more than twice in HEK293^{pSsiPKD1} cells by NCCE inhibitors, but remains unchanged in HEK293^{pSUPER} cells. Results have been obtained in two independent experiments and expressed as means \pm SD.

PC1 expression by RNA interference in HEK293 cells or loss of PC1 expression in ADPKD kidney cystic cells increases oscillatory activity of cytoplasmic Ca^{2+} , evoked by low serum concentrations. Consistently, we have also found that expression of exogenous PC1 reduces the frequency of serum-evoked Ca^{2+} oscillations not only in PC1-deficient HEK293^{pSsiPKD1} cells, but also in HEK293 control cells. The main conclusion of this study is therefore that PC1 is required for negative control of Ca^{2+} oscillation in kidney-derived cells. The implication is that PC1 exerts a profound influence on Ca^{2+} signalling by controlling Ca^{2+} oscillation induced by serum, with a crucial role in cell proliferation. Cell growth studies have shown that cell proliferation in PC1-deficient HEK293 cells is inversely related to the level of PC1 expression which, in turn, is inversely related to the Ca^{2+} oscillation frequency.

Ca²⁺ oscillation and cell proliferation

Intracellular Ca²⁺ homeostasis appears to be altered in HEK293^{pSsiPKD1} cells with basal Ca²⁺ levels higher than those of controls. Increased Ca²⁺ levels have already been observed in the HEK293 cells expressing the dominant negative PC1 C-terminus (Puri *et al.* 2004; Manzati *et al.* 2005). These results therefore indicate that loss of PC1 function may be associated with increase in basal Ca²⁺ levels, which may contribute to increased proliferation of these cells. This hypothesis is also supported by the observation that overexpression of TRPV6 slightly increases intracellular Ca²⁺ levels and Ca²⁺-dependent proliferation of HEK293 cells (Schwarz *et al.* 2006). However, in primary cultures of ADPKD cystic cells reduction in basal Ca²⁺ levels has been documented, imputing abnormal proliferation of cystic cells to this alteration (Yamaguchi *et al.* 2006). Even though characterization of molecular lesions in primary ADPKD cultures has not been reported, these differences in basal Ca²⁺ levels may be related to the different type of cell culture. Nevertheless, in the light of these contrasting results it is difficult to explain the PKD-dependent abnormal proliferation with abnormal basal Ca²⁺ levels.

Interestingly, in HEK293^{pSsiPKD1} cells, serum treatment that increased cell proliferation, did not produce a significant increase in evoked Ca²⁺ levels (Fig. 2a), but rather a clear change (2-fold statistically significant increase) in frequency of FBS-induced Ca²⁺ oscillations. Notably, frequency of Ca²⁺ oscillation is also increased in cystic cells either heterozygous or hemizygous for the Q2556X PKD1. Because both cystic cell lines show increased FBS-evoked Ca²⁺ oscillations, these may be a cause of aberrant proliferation in these cells as well as in HEK293^{pSsiPKD1} cells. It must, however, be considered that the present study was performed in immortalized cells. Although these are derived from normal and cyst-lining epithelia and maintain many features of tubule of origin, immortalized cells are poorly differentiated and polarized, particularly when cultured under subconfluent conditions such as those used for a proliferation study. Therefore, the precise role of abnormal Ca²⁺ oscillations on cyst formation and cell population growth in the kidney remains to be ascertained.

The finding that HEK293 cells expressing mouse PC1 showed reductions in both Ca²⁺ oscillation and cell proliferation, provides evidence for the role of Ca²⁺ oscillations in the PC1-dependent control of cell proliferation. In the light of these observations therefore growth suppression activity of PC1 may be explained by its inhibitory action on Ca²⁺ oscillations as was observed in both normal and HEK293^{pSsiPKD1} cells exogenously expressing the mouse PC1.

Ca²⁺ channel activity and cell proliferation

Ca²⁺ entry that supports [Ca²⁺]_i cycling, predominantly occurs through store-operated or a CCE pathway (Bird & Putney 2005). Nevertheless, in many cell types, including HEK293 cells, extracellular Ca²⁺ entry and Ca²⁺ oscillations also result from NCCE, whose activation is independent of store depletion (Mignen *et al.* 2001).

Non-capacitative Ca²⁺ entry activity appears to be crucial for HEK293^{pSsiPKD1} cell proliferation. Ca²⁺ oscillations in HEK293^{pSsiPKD1} cells are insensitive to 1 μM Gd³⁺, but markedly inhibited by the NCCE blocker, which also inhibited cell proliferation. Moreover, in the absence of external Ca²⁺, no difference was observed in 1% evoked Ca²⁺ oscillations between HEK293^{pSsiPKD1} and HEK293^{pSUPER} control cells (3.34 ± 0.31 versus 2.58 ± 0.9 oscillations in 15 min/cell, respectively), consistently with a role of NCCE activity on the oscillation increase. Therefore, the PC1 loss leads to deregulation of the NCCE pathway, strongly associating increased Ca²⁺ oscillations and proliferation to abnormal activity of NCCE channels. Consistently, both SKF96365 and LOE908 are able to reduce proliferation in PKD1-mutated heterozygous and hemizygous cystic cells, but not wild-type kidney tubule cells. Ca²⁺ influx through NCCE channels is therefore necessary for the increased [Ca²⁺]_i oscillatory response and cell proliferation in PC1-deficient cells.

On the basis of these considerations, we hypothesize that PC1 controls an as yet uninvestigated activity of NCCE channels. In HEK293 cells, NCCE has been shown to be associated to the activity of the arachidonate-regulated Ca²⁺ channel (ARC channel) (Mignen *et al.* 2003a), but because we have observed that calcineurin activity is required for the FBS-induced proliferation increase in HEK293^{pSsiPKD1} cells (Fig. 7d), and because ARC channels are turned off by calcineurin activation (Mignen *et al.* 2003b), these channels do not seem to be the target of the PC1 control. On the other hand, an arachidonate-activated NCCE channel activity, which displays biophysical properties different from the ARC channels of HEK293 cells, has been reported in endothelial cells (Fiorio Pla & Munaron 2001). Interestingly, this NCCE activity is required for the promotion of cell proliferation activated by peptidic growth factors like the basic fibroblast growth factor (Fiorio Pla & Munaron 2001). This NCCE activity is also involved in peripheral and localized Ca²⁺ events, probably leading to the involvement of differential patterns of Ca²⁺-dependent proteins and genes (Tomatis *et al.* 2007), including calcineurin and PKC pathways, abnormally activated in HEK293^{pSsiPKD1} cells. This activity may lead, in addition to increased cell proliferation, to the abnormal polarity, migration and apoptosis typically observed in ADPKD cystic cells (Ong & Harris 2005).

Ca²⁺ oscillations and signalling

The results of our studies suggest that a loss of PC1 in HEK293^{pSsiPKD1} cells promotes a serum-induced proliferation increase by inducing Ca²⁺ entry and subsequent activation of the calcineurin/NFAT pathway. As reported in other cell types (Lipskaia & Lompre 2004), NFAT activation in HEK293^{pSsiPKD1} cells promotes cell cycle progression in association with a reduction in the activity of cyclin kinase inhibitors. Consistently, cyclosporin A treatment, which, by inhibiting calcineurin, inhibits NFAT dephosphorylation and its nuclear translocation (Hogan *et al.* 2003), blocks the PC1-dependent increase in proliferation (Fig. 7d). Here, we also show that once NCCE-mediated Ca²⁺ entry is blocked by SKF96365 treatment in HEK293^{pSsiPKD1} cells, NFAT activation is markedly reduced. Thus, as previously demonstrated in other cell types (Dolmetsch *et al.* 1998; Tomida *et al.* 2003), activation of NFAT in HEK293^{pSsiPKD1} appears to be linked to increase in Ca²⁺ oscillation, possibly accompanied by the release of Ca²⁺ from intracellular stores and activation of store-operated Ca²⁺ entry, as proposed in cells expressing the PC1 C-terminus (Puri *et al.* 2004). Because this peptide appears to act as a dominant negative (Manzati *et al.* 2005; Xu *et al.* 2007), NFAT activation in these cells is consistent with that found in HEK^{pSsiPKD1} cells. HEK293^{pSsiPKD1} cell proliferation requires PKC- α activation and p27 down-regulation (see Fig. 8), as in cells expressing the PC1 C-terminus that show reduced levels of both p27^{Kip} and p21^{Waf} associated to increased cell proliferation (Manzati *et al.* 2005). This observation is consistent with previous findings showing that exogenous expression of full-length PC1 in kidney cells induces cell cycle arrest in G₀/G₁ by up-regulating p21^{Waf} expression through the JAK-STAT pathway (Bhunia *et al.* 2002). It has also been reported that calcineurin and NFAT activity is required in keratinocytes for the expression of p21 and p27, which contribute to growth arrest of these cells (Santini *et al.* 2001). This supports the postulated cross-talk between calcineurin, p21 and p27. In HEK293^{pSsiPKD1}, this cross-talk may either be dysfunctional or p27 gene down-regulation may result by one or more different Ca²⁺-dependent signalling pathways activated by PC1 depletion (see Fig. 8).

Furthermore, the association between PKC- α membrane translocation and Ca²⁺ oscillation is consistent with the activation of PKC by glutamate-induced Ca²⁺ waves in astrocytes (Codazzi *et al.* 2001).

Because NFAT nuclear translocation occurs in HEK293^{pSsiPKD1} and control cells only after serum treatment, the role of basal Ca²⁺ again appears to be irrelevant in the proliferation increase of




Figure 9. A hypothetical model illustrating the role of PC1 on the agonist-evoked NCCE channel activity and effects of Ca²⁺ oscillations (Ca²⁺Os) on NFAT pathway and cell proliferation. In HEK293^{pSUPER} and tubular kidney control cells, PC1 negatively modulates NCCE activity through a still undefined direct or indirect mechanism (dotted bar); Ca²⁺ oscillations are dependent on both CCE and NCCE activities as shown by comparable inhibitory effects of 1 μM Gd³⁺ and SKF96365 on Ca²⁺ oscillations and cell proliferation (Figs 5 and 6). Loss of PC1 function causes an increase in NCCE channel activity and in Ca²⁺ oscillations thus leading to an increase in Ca²⁺-dependent NFAT activation and cell proliferation.

PC1-deficient cells, while the role of Ca²⁺ oscillations appears to be crucial. These results provide new insight into the molecular and physiological mechanism of Ca²⁺ oscillations in PKD1-like kidney cells and in promoting NFAT-dependent kidney cell proliferation.

In summary, our results establish that the molecular suppression of endogenous PC1 in HEK293 cells and PKD1 cystic cells leads to abnormal Ca²⁺ oscillations and has a profound influence on Ca²⁺ signalling and cell proliferation, as illustrated in Fig. 9. At present, it is impossible to say how many steps may lie between PC1 and the NCCE channel. Because PC1 has been implicated in gene expression control by activating a variety of transcription factors including the NFAT investigated here, NCCE modulation may result from changes in gene expression, or in post-translational modification (Li *et al.* 2005). Moreover, either a direct or indirect interaction between PC1 and NCCE channel may exist. PC1 may, in fact, constitute Ca²⁺-permeable channel activities either by interacting with PC2 in the PC1/PC2 receptor-ion channel (Hanaoka *et al.* 2000), or independently of PC2 (Babich *et al.* 2004).

In conclusion, our findings suggest that PC1 may function as a controller of Ca²⁺ release in response to extracellular stimuli also through the NCCE pathway, thus regulating the Ca²⁺-mediated signal transduction involved in cell proliferation through the NFAT pathway. Moreover, because NCCE channel blockers may reduce the PC1 suppression-dependent [Ca²⁺]_i oscillation as well

as cell proliferation, these channels could be potential therapeutic targets for treatment of the abnormalities observed in cystic cells.

ACKNOWLEDGEMENTS

We thank Dr. A.L. Armesilla (University of Manchester) and Dr. E. Olson (South-western University, Dallas, TX, USA) for providing NFAT promoter-luciferase reporter and GFP-tagged NFAT2 plasmids, respectively, Consugar M. (Mayo Clinic, Rochester, MN, USA) for PKD1-mutated cell line cultures, and Anna Foster for correcting the manuscript. LOE908 was a kind gift from Boehringer Ingelheim Pharma GmbH & Co. KG (Biberach, Germany). This work was supported by grants from Italian Telethon GTF01011, Italian MIUR COFIN 2002, FIRB 2005, Cassa di Risparmio di Ferrara, Local University Funds, and the Italian Association for Cancer Research.

REFERENCES

- Aguiari G, Piva R, Manzati E, Mazzoni E, Augello G, Chiari E, Moretti S, Neri LM, del Senno L (1998) K562 erythroid and HL60 macrophage differentiation downregulates polycystin, a large membrane-associated protein. *Exp. Cell Res.* **244**, 259–267.
- Aguiari G, Campanella M, Manzati E, Pinton P, Banzi M, Moretti S, Piva R, Rizzuto R, del Senno L (2003) Expression of polycystin-1 C-terminal fragment enhances the ATP-induced Ca²⁺ release in human kidney cells. *Biochem. Biophys. Res. Commun.* **301**, 657–664.
- Aguiari G, Banzi M, Gessi S, Cai Y, Zeggio E, Manzati E, Piva R, Lambertini E, Ferrari L, Peters DJ, Lanza F, Harris PC, Borea PA, Somlo S, Del Senno L (2004) Deficiency of polycystin-2 reduces Ca²⁺ channel activity and cell proliferation in ADPKD lymphoblastoid cells. *FASEB J.* **18**, 884–886.
- Babich V, Zeng WZ, Yeh BI, Ibraghimov-Beskrovnaya O, Cai Y, Somlo S, Huang CL (2004) The N-terminal extracellular domain is required for polycystin-1-dependent channel activity. *J. Biol. Chem.* **279**, 25582–25589.
- Berridge MJ (2002) The endoplasmic reticulum: a multifunctional signaling organelle. *Cell Calcium* **32**, 235–249.
- Berridge MJ, Lipp P, Bootman MD (2000) The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell Biol.* **1**, 11–21.
- Bhunia AK, Piontek K, Boletta A, Liu L, Qian F, Xu PN, Germino FJ, Germino GG (2002) PKD1 induces p21 (waf1) and regulation of the cell cycle via direct activation of the JAK-STAT signaling pathway in a process requiring PKD2. *Cell* **109**, 157–168.
- Bird GS, Putney JW Jr (2005) Capacitative calcium entry supports calcium oscillations in human embryonic kidney cells. *J. Physiol.* **562**, 697–706.
- Boca M, Distefano G, Qian F, Bhunia AK, Germino GG, Boletta A (2006) Polycystin-1 induces resistance to apoptosis through the phosphatidylinositol 3-kinase/Akt signaling pathway. *J. Am. Soc. Nephrol.* **17**, 637–647.
- Brummelkamp TR, Bernards R, Agami R (2002) A system for stable expression of short interfering RNAs in mammalian cells. *Science* **296**, 550–553.
- Codazzi F, Teruel MN, Meyer T (2001) Control of astrocyte Ca²⁺ oscillations and waves by oscillating translocation and activation of protein kinase C. *Curr. Biol.* **11**, 1089–1097.
- Delmas P (2004) Polycystins: from mechanosensation to gene regulation. *Cell* **118**, 145–148.
- Delmas P, Nomura H, Li X, Lakkis M, Luo Y, Segal Y, Fernandez-Fernandez JM, Harris P, Frischauf AM, Brown DA, Zhou J (2002) Constitutive activation of G-proteins by polycystin-1 is antagonized by polycystin-2. *J. Biol. Chem.* **277**, 11276–11283.
- Dolmetsch RE, Xu K, Lewis RS (1998) Calcium oscillations increase the efficiency and specificity of gene expression. *Nature* **392**, 933–936.
- Fiorio Pla A, Munaron L (2001) Calcium influx, arachidonic acid, and control of endothelial cell proliferation. *Cell Calcium* **30**, 235–244.

- Gabow PA (1990) Autosomal dominant polycystic kidney disease – more than a renal disease. *Am. J. Kidney Dis.* **16**, 403–413.
- González-Perrett S, Kim K, Ibarra C, Damiano AE, Zotta E, Batelli M, Harris PC, Reisin IL, Arnaout MA, Cantiello HF (2001) Polycystin-2, the protein mutated in autosomal dominant polycystic kidney disease (ADPKD), is a Ca²⁺-permeable nonselective cation channel. *Proc. Natl. Acad. Sci. USA* **98**, 1182–1187.
- Grimm DH, Cai Y, Chauvet V, Rajendran V, Zeltner R, Geng L, Avner ED, Sweeney W, Somlo S, Caplan MJ (2003) Polycystin-1 distribution is modulated by polycystin-2 expression in mammalian cells. *J. Biol. Chem.* **278**, 36786–36793.
- Hanaoka K, Qian F, Boletta A, Bhunia AK, Piontek K, Tsiokas L, Sukhatme VP, Guggino WB, Germino GG (2000) Co-assembly of polycystin-1 and -2 produces unique cation-permeable currents. *Nature* **408**, 990–994.
- Harris PC (1999) Autosomal dominant polycystic kidney disease: clues to pathogenesis. *Hum. Mol. Genet.* **8**, 1861–1866.
- Hogan PG, Chen L, Nardone J, Rao A (2003) Transcriptional regulation by calcium, calcineurin, and NFAT. *Genes Dev.* **17**, 2205–2232.
- Kim H, Bae Y, Jeong W, Ahn C, Kang S (2004) Depletion of PKD1 by antisense oligodeoxynucleotide induces premature G₁/S-phase transition. *Eur. J. Hum. Genet.* **12**, 433–440.
- Koulen P, Cai Y, Geng L, Maeda Y, Nishimura S, Witzgall R, Ehrlich BE, Somlo S (2002) Polycystin-2 is an intracellular calcium release channel. *Nat. Cell Biol.* **4**, 191–197.
- Li X, Luo Y, Starremans PG, McNamara CA, Pei Y, Zhou J (2005) Polycystin-1 and polycystin-2 regulate the cell cycle through the helix-loop-helix inhibitor Id2. *Nat. Cell Biol.* **7**, 1202–1212.
- Lipskaia L, Lompre AM (2004) Alteration in temporal kinetics of Ca²⁺ signaling and control of growth and proliferation. *Biol. Cell* **96**, 55–68.
- Manzati E, Aguiari G, Banzi M, Manzati M, Selvatici R, Falzarano S, Maestri I, Pinton P, Rizzuto R, del Senno L (2005) The cytoplasmic C-terminus of polycystin-1 increases cell proliferation in kidney epithelial cells through serum-activated and Ca²⁺-dependent pathway(s). *Exp. Cell Res.* **304**, 391–406.
- Mignen O, Thompson JL, Shuttleworth TJ (2001) Reciprocal regulation of capacitative and arachidonate-regulated non-capacitative Ca²⁺ entry pathways. *J. Biol. Chem.* **276**, 35676–35683.
- Mignen O, Thompson JL, Shuttleworth TJ (2003a) Ca²⁺ selectivity and fatty acid specificity of the noncapacitative, arachidonate-regulated Ca²⁺ (ARC) channels. *J. Biol. Chem.* **278**, 10174–10181.
- Mignen O, Thompson JL, Shuttleworth TJ (2003b) Calcineurin directs the reciprocal regulation of calcium entry pathways in nonexcitable cells. *J. Biol. Chem.* **278**, 40088–40096.
- Mochizuki T, Wu G, Hayashi T, Xenophontos SL, Veldhuisen B, Saris JJ, Reynolds DM, Cai Y, Gabow PA, Pierides A, Kimberling WJ, Breuning MH, Deltas CC, Peters DJ, Somlo S (1996) PKD2, a gene for polycystic kidney disease that encodes an integral membrane protein. *Science* **272**, 1339–1342.
- Moneer Z, Taylor CW (2002) Reciprocal regulation of capacitative and non-capacitative Ca²⁺ entry in A7r5 vascular smooth muscle cells: only the latter operates during receptor activation. *Biochem. J.* **362**, 13–21.
- Moneer Z, Dyer JL, Taylor CW (2003) Nitric oxide co-ordinates the activities of the capacitative and non-capacitative Ca²⁺-entry pathways regulated by vasopressin. *Biochem. J.* **370**, 439–448.
- Nauli SM, Rossetti S, Kolb RJ, Alenghat FJ, Consugar MB, Harris PC, Ingber DE, Loghman-Adham M, Zhou J (2006) Loss of polycystin-1 in human cyst-lining epithelia leads to ciliary dysfunction. *J. Am. Soc. Nephrol.* **17**, 1015–1025.
- Nickel C, Benzing T, Sellin L, Gerke P, Karihaloo A, Liu ZX, Cantley LG, Walz G (2002) The polycystin-1 C-terminal fragment triggers branching morphogenesis and migration of tubular kidney epithelial cells. *J. Clin. Invest.* **109**, 481–489.
- Ong AC, Harris PC (2005) Molecular pathogenesis of ADPKD: the polycystin complex gets complex. *Kidney Int.* **67**, 1234–1247.
- Parnell SC, Magenheimer BS, Maser RL, Rankin CA, Smine A, Okamoto T, Calvet JP (1998) The polycystic kidney disease-1 protein, polycystin-1, binds and activates heterotrimeric G-proteins *in vitro*. *Biochem. Biophys. Res. Commun.* **251**, 625–631.
- Pelucchi B, Aguiari G, Pignatelli A, Manzati E, Witzgall R, Del Senno L, Belluzzi O (2006) Nonspecific cation current associated with native polycystin-2 in HEK-293 cells. *J. Am. Soc. Nephrol.* **17**, 388–397.
- Polgar K, Burrow CR, Hyink DP, Fernandez H, Thornton K, Li X, Gusella GL, Wilson PD (2005) Disruption of polycystin-1 function interferes with branching morphogenesis of the ureteric bud in developing mouse kidneys. *Dev. Biol.* **286**, 16–30.
- Puri S, Magenheimer BS, Maser RL, Ryan EM, Zien CA, Walker DD, Wallace DP, Hempson SJ, Calvet JP (2004) Polycystin-1 activates the calcineurin/NFAT (nuclear factor of activated T-cells) signaling pathway. *J. Biol. Chem.* **279**, 55455–55464.
- Qian Q, Li M, Cai Y, Ward CJ, Somlo S, Harris PC, Torres VE (2003) Analysis of the polycystins in aortic vascular smooth muscle cells. *J. Am. Soc. Nephrol.* **14**, 2280–2287.

- Santini MP, Talora C, Seki T, Bolgan L, Dotto GP (2001) Cross talk among calcineurin, Sp1/Sp3, and NFAT in control of p21 (WAF1/CIP1) expression in keratinocyte differentiation. *Proc. Natl. Acad. Sci. USA* **98**, 9575–9580.
- Schwarz EC, Wissenbach U, Niemeyer BA, Strauß B, Philipp SE, Flockerzi V, Hoth M (2006) TRPV6 potentiates calcium-dependent cell proliferation. *Cell Calcium* **39**, 163–173.
- Tomatis C, Fiorio Pla A, Munaron L (2007) Cytosolic calcium microdomains by arachidonic acid and nitric oxide in endothelial cells. *Cell Calcium* **41**, 261–269.
- Tomida T, Hirose K, Takizawa A, Shibasaki F, Iino M (2003) NFAT functions as a working memory of Ca²⁺ signals in decoding Ca²⁺ oscillation. *EMBO J.* **22**, 3825–3832.
- Vandorpe DH, Chernova MN, Jiang L, Sellin LK, Wilhelm S, Stuart-Tilley AK, Walz G, Alper SL (2001) The cytoplasmic C-terminal fragment of polycystin-1 regulates a Ca²⁺-permeable cation channel. *J. Biol. Chem.* **276**, 4093–4101.
- Xu C, Rossetti S, Jiang L, Harris PC, Brown-Glaberman U, Wandering-Ness A, Bacallao R, Alper SL (2007) Human ADPKD primary cyst epithelial cells with a novel, single codon deletion in the PKD1 gene exhibit defective ciliary polycystin localization and loss of flow-induced Ca²⁺ signaling. *Am. J. Physiol. Renal Physiol.* **292**, 930–945.
- Yamaguchi T, Hempson SJ, Reif GA, Hedge AM, Wallace DP (2006) Calcium restores a normal proliferation phenotype in human polycystic kidney disease epithelial cells. *J. Am. Soc. Nephrol.* **17**, 178–187.